

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 December 2002 (12.12.2002)

PCT

(10) International Publication Number
WO 02/099079 A2

(51) International Patent Classification⁷: C12N

(74) Agent: DECAMP, James, D.; Clark & Elbing LLP, 101
Federal Street, Boston, MA 02110 (US).

(21) International Application Number: PCT/US02/18066

(22) International Filing Date: 6 June 2002 (06.06.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/296,554 6 June 2001 (06.06.2001) US

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 60/296,554 (CIP)
Filed on 6 June 2001 (06.06.2001)

(71) Applicant (for all designated States except US): THE
GENERAL HOSPITAL CORPORATION [US/US]; 55
Fruit Street, Boston, MA 02114 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SHEEN, Jen
[US/US]; 9 Hawthorne Place, Apartment 5J, Boston, MA
02114 (US). HWANG, Ildoo [KR/US]; 29J Montrose
Manor Court, Baltimore, MD 21228 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: CYTOKININ RESPONSE REGULATORS AND USES THEREOF

(57) Abstract: The invention generally features methods for increasing yield, shoot formation, and delaying senescence in plants with the use of transgenes that regulate the cytokinin response. The invention also features plants and plant components that harbor the transgene(s).



WO 02/099079 A2

CYTOKININ RESPONSE REGULATORS AND USES THEREOF

5

Background of the Invention

This invention relates to genetically-engineered plants having increased yield and productivity, shoot, leaf and meristem formation, enhanced photosynthesis, and delayed senescence.

10 Despite long recognition of cytokinins as essential plant hormones involved in diverse processes of plant growth and development, including cell division, shoot initiation, leaf and root differentiation, chloroplast biogenesis, apical dominance, and senescence, the molecular and biochemical mechanisms underlying cytokinin actions have not been elucidated (Davies, *Plant Hormones: Physiology, Biochemistry and Molecular Biology*, 1995, Kluwer Academic Publishers, Dordrecht; Meijer and Murray, *Curr. Opin. Plant Biol.* 4: 44-9, 2001; Mok and Mok, *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 52: 89-118, 2001; Quirino et al., *Trends Plant Sci.* 5: 278-82, 2000). Recent genetic identification of *Arabidopsis* hybrid histidine protein kinases (AHKs), CKI1 and CRE1, in
15 cytokinin signalling (Kakimoto, *Science* 274: 982-5, 1996; Suzuki et al., *Plant Cell. Physiol.* 42: 107-13, 2001; Inoue et al., *Nature* 409: 1060-3, 2001), the characterization of *Arabidopsis* response regulators (ARRs) as cytokinin primary response genes (D'Agostino et al., *Plant Physiol.* 124: 1706-17, 2000; Kiba et al., *Plant Cell Physiol.* 40: 767-71, 1999), and the existence of *Arabidopsis* histidine-
20 containing phosphotransmitters (AHPs; Suzuki, Imamura, et al., *Plant Cell Physiol.* 39: 1258-68, 1998), have implicated the involvement of a two-component phosphorelay mechanism in cytokinin signal transduction. However, cellular and molecular evidence is lacking in supporting a complete cytokinin signalling circuit in plant cells.

30 Two-component circuitry, consisting of a histidine kinase (HK) sensor and a response regulator (RR) output, are responsible for signal transduction in most prokaryotic and some eukaryotic systems. The signalling pathway is initiated by

a HK sensor and mediated by phosphotransfer between a conserved histidine (His) residue in HKs or histidine-containing phosphotransmitters (HPs) and a conserved aspartate (Asp) residue in RRs (Wurgler-Murphy and Saito, *Trends Biochem. Sci.* 22: 172-6, 1997; Stock et al., *Annu. Rev. Biochem.* 69: 183-215, 2000). Since there are only one HK, one HP, and two RRs in *S. cerevisiae*, it was previously believed that two-component signal transduction has limited function in eukaryotes (Stock et al., *supra*). However, the completion of the *Arabidopsis* genome has revealed over 40 genes encoding putative two-component signal transducers, AHKs, AHPs and ARRs, suggesting a significant involvement of the ancient and conserved signalling mechanism in many facets of plant cell regulation (Urao, et al., *Trends Plant Sci.* 5: 67-74, 2000). The identification of conserved HK signature motifs and/or activity in the photoreceptor phytochrome, putative osmosensor, and the ethylene and cytokinin receptors in *Arabidopsis* further supports this view (Inoue et al., *supra*; Urao, et al., *supra*; Yeh and Lagarias, *Proc. Natl. Acad. Sci. U S A* 95: 13976-81, 1998; Bleecker and Kende, *Annu. Rev. Cell Dev. Biol.* 16: 1-18, 2000).

In eukaryotic signal transduction, the two-component circuit often provides a link between the HK sensor to a MAP kinase (MAPK) signaling cascade. For example, the osmosensing signal transduction pathway in yeast is mediated by the SLN1/YPD1/SSK1 phosphorelay. The RR SSK1 then activates the HOG1 MAPK cascade in the cytosol to control gene expression (Maeda et al., *Nature* 369: 242-5, 1994; Posas et al., *Cell* 86: 865-75, 1996; Posas and Saito, *EMBO J.* 17: 1385-94, 1998). It has been speculated that plant HKs such as the ethylene receptor (ETR1) and a putative osmosensor (AHK1) can initiate a phosphorelay and transmit the signal through a MAPK cascade (Bleecker and Kende, *supra*; Gamble et al., *Proc. Natl. Acad. Sci. U S A* 95: 7825-9, 1998; Clark et al., *Proc. Natl. Acad. Sci. U S A* 95: 5401-6, 1998; Urao et al., *Plant Cell* 11: 1743-54, 1999). Due to the lack of physiological plant cell assays, the mechanisms of HK action and the signal transduction pathways of any HK-mediated plant responses, including cytokinin signalling, remain obscure.

Summary of the Invention

The present invention capitalizes on the discovery that manipulation of the expression of cytokinin response regulators increases plant yield growth, and
5 productivity.

In one aspect, the invention features a method for increasing yield in a plant, the method including the steps of: (a) introducing into plant cells a transgene including DNA encoding a B-type response regulator operably linked to a promoter functional in plant cells to yield transformed plant cells; and (b)
10 regenerating a plant from the transformed cells, wherein the B-type response regulator is expressed in the cells of the transgenic plant, thereby increasing yield in the plant. In preferred embodiments, the B-type response regulator is a crucifer B-type response regulator (for example, ARR1 (SEQ ID NO.: 2); ARR2 (SEQ ID NO.: 3); and ARR10 (SEQ ID NO.: 9)).

15 In a second aspect, the invention features a method for increasing yield in a plant, the method including the steps of: (a) introducing into plant cells a transgene operably linked to a promoter functional in plant cells to yield transformed plant cells; and (b) regenerating a plant from the transformed cells, wherein expression of the transgene reduces expression of an A-type response
20 regulator in the cells of the plant, thereby increasing yield in the plant. In preferred embodiments, the A-type response regulator is a crucifer A-type response regulator (for example, ARR 4 (SEQ ID NO.: 5); ARR 5 (SEQ ID NO.: 6); ARR 6 (SEQ ID NO.: 7); and ARR 7 (SEQ ID NO.: 8)). In other preferred embodiments, the transgene expresses an antisense molecule of A-type response
25 regulator; a dominant negative gene product of A-type response regulator; or expression of the transgene results in the co-suppression of the A-type response regulator.

In a third aspect, the invention features a method for increasing shoot formation in a plant, the method including the steps of: (a) introducing into plant
30 cells a transgene including DNA encoding a B-type response regulator operably

linked to a promoter functional in plant cells to yield transformed plant cells; and
(b) regenerating a plant from the transformed cells, wherein the B-type response
regulator is expressed in the cells of the plant, thereby increasing shoot formation
in the plant. In preferred embodiments, the method includes the use of a B-type
5 response regulator that is a crucifer B-type response regulator.

In a fourth aspect, the invention features a method for increasing shoot
formation in a plant, the method including the steps of: (a) introducing into plant
cells a transgene operably linked to a promoter functional in plant cells to yield
transformed plant cells; and (b) regenerating a plant from the transformed cells,
10 wherein expression of the transgene reduces expression of an A-type response
regulator in the cells of the plant, thereby increasing shoot formation in the plant.

In a fifth aspect, the invention features a method for delaying senescence
in a plant, the method including the steps of: (a) introducing into plant cells a
transgene including DNA encoding a B-type response regulator operably linked
15 to a promoter functional in plant cells to yield transformed plant cells; and (b)
regenerating a plant from the transformed cells, wherein the B-type response
regulator is expressed in the cells of the plant, thereby delaying senescence in the
plant.

In a sixth aspect, a method for delaying senescence in a plant, the method
20 including the steps of: (a) introducing into plant cells a transgene operably linked
to a promoter functional in plant cells to yield transformed plant cells; and (b)
regenerating a plant from the transformed cells, wherein expression of the
transgene reduces expression of an A-type response regulator in the cells of the
plant, thereby delaying senescence in the plant.

25 In a seventh aspect, the invention features a method for increasing yield in
a plant, the method including the steps of: (a) introducing into plant cells a
transgene including DNA encoding a histidine kinase operably linked to a
promoter functional in plant cells to yield transformed plant cells; and (b)
regenerating a plant from the transformed cells, wherein histidine kinase is

expressed in the cells of the plant, thereby increasing yield in the plant. In preferred embodiments, the histidine kinase is a crucifer histidine kinase (for example, CKI1 (SEQ ID NO.: 13) and CRE1 (SEQ ID NO.: 18)).

In an eighth aspect, the invention features a method for increasing shoot
5 formation in a plant, the method including the steps of: (a) introducing into plant cells a transgene including DNA encoding a histidine kinase operably linked to a promoter functional in plant cells to yield transformed plant cells; and (b) regenerating a plant from the transformed cells, wherein histidine kinase is expressed in the cells of the plant, thereby increasing shoot formation in the plant.

10 In a ninth aspect, a method for delaying senescence in a plant, the method including the steps of: (a) introducing into plant cells a transgene including DNA encoding a histidine kinase operably linked to a promoter functional in plant cells to yield transformed plant cells; and (b) regenerating a plant from the transformed cells, wherein histidine kinase is expressed in the cells of the plant, thereby
15 delaying senescence in the plant.

The invention further includes plants and plant components expressing one, or a combination of two or more, of the aforementioned transgenes. The invention also encompasses methods of generating these plants or plant components, which may involve introducing the transgenes individually to
20 separate plants or plant components, and then crossing the appropriate genotypes to get the desired transgene combination. Alternatively, one or more constructs expressing a desired combination of transgenes may be generated and then introduced into the plant or plant component.

For example, the invention also relates to a plant or plant component
25 comprising at least one transgene encoding (i) an A-type response regulator polypeptide, (ii) an antisense A-type response regulator RNA, (iii) a dominant negative A-type response regulator polypeptide, (iv) a B-type response regulator polypeptide, (v) an antisense B-type response regulator RNA, (vi) a dominant negative B-type response regulator, (vii) an antisense HK RNA, (viii) a dominant

30

negative HK polypeptide, or any combination of (i) –(viii). In addition, the invention includes any of the aforementioned plants further including a transgene encoding a histidine kinase polypeptide.

The invention further includes a plant or plant component that includes:

- 5 (a) a first transgene encoding (i) an A-type response regulator polypeptide, (ii) an antisense A-type response regulator RNA, or (iii) a dominant negative A-type response regulator polypeptide or combination thereof; (b) a second transgene encoding (iv) a B-type response regulator polypeptide, (v) an antisense B-type response regulator RNA, (vi) a dominant negative B-type response regulator or
10 combination thereof, and (c) a third transgene encoding (vii) a HK polypeptide; (viii) an antisense HK RNA, (ix) a dominant negative HK polypeptide or combination thereof.

Exemplary plants which are useful in the methods of the invention, as well as for generating the plants (or plant cells, plant components, plant tissues, or
15 plant organs) of the invention, include dicots and monocots, such as sugar cane, wheat, rice, maize, sugar beet, barley, manioc, crucifer, mustard, potato, soybean, sorghum, cassava, banana, grape, oats, tomato, millet, coconut, orange, rye, cabbage, apple, eggplant, watermelon, canola, cotton, carrot, garlic, onion, pepper, strawberry, yam, papaya, peanut, onion, legume, bean, pea, mango, and
20 sunflower.

By “operably linked” is meant that a gene and a regulatory sequence(s) are connected in such a way as to permit gene expression when the appropriate molecules (for example, transcriptional activator proteins) are bound to the regulatory sequence(s).

25 By “plant cell” is meant any self-propagating cell bounded by a semi-plant component expression control permeable membrane and containing a plastid. Such a cell also requires a cell wall if further propagation is desired. Plant cell, as used herein, includes, without limitation, algae, cyanobacteria, seeds, suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots,
30 gametophytes, sporophytes, pollen, and microspores.

By "plant component" is meant a part, segment, or organ obtained from an intact plant or plant cell. Exemplary plant components include, without limitation, somatic embryos, leaves, stems, roots, flowers, tendrils, fruits, scions, and rootstocks.

5 By "transgene" is meant any piece of DNA which is inserted by artifice into a cell, and becomes part of the genome of the organism which develops from that cell. Such a transgene may include a gene which is partly or entirely heterologous (i.e., foreign) to the transgenic organism, or may represent a gene homologous to an endogenous gene of the organism.

10 By "transgenic" is meant any cell which includes a nucleic acid sequence (e.g., a recombinant DNA sequence) which is inserted by artifice into a cell and becomes part of the genome of the organism which develops from that cell. As used herein, the transgenic organisms are generally transgenic plants and the DNA (transgene) is inserted by artifice into the nuclear or plastidic genome.

15 By "yield" or "plant yield" is meant increased growth (e.g., crop growth) or increased biomass. For example, increased yield results from increased shoot growth or meristem formation. Plants expressing the genes disclosed herein exhibiting increased yield can be selected by visual observation, for example by comparison with a wild-type plant.

20 By "reducing expression" or "reduces expression" is meant a decrease in the level of gene expression (for example, expression of a gene encoding a A-type response regulator) by at least 30-50%, preferably by at least 50-80%, and more preferably by at least 80-95% or greater relative to the level in a control plant (for example, a wild-type plant). Reduction of such expression levels may be
25 accomplished by employing standard methods which are known in the art including, without limitation, antisense and co-suppression technologies, expression of a dominant negative gene product, or through the generation of mutated genes using standard mutagenesis techniques. Levels of negative

regulator polypeptide or transcript are monitored according to any standard technique including, but not limited to, northern blotting, RNase protection, or immunoblotting.

By "crucifer" is meant any plant that is classified within the Cruciferae family. The Cruciferae include many agricultural crops, including, without limitation, rape (for example, *Brassica campestris* and *Brassica napus*), broccoli, cabbage, brussel sprouts, radish, kale, Chinese kale, kohlrabi, cauliflower, turnip, rutabaga, mustard, horseradish, and *Arabidopsis*.

By "a promoter functional in a plant cell" is meant any minimal sequence sufficient to direct transcription in a plant cell. Included in the invention are promoter elements that are sufficient to render promoter-dependent gene expression controllable for cell-, tissue-, or organ-specific gene expression, or elements that are inducible by external or internal agents (for example, cytokinins), or elements that are capable of cycling gene transcription; such elements may be located in the 5' or 3' regions of the native gene or engineered into a transgene construct.

The invention provides a number of important advances and advantages for improving and enhancing agronomically important traits such as photosynthesis, productivity, yield, leaf, shoot, and meristem formation, and delaying senescence. In particular, the invention provides for increased production efficiency, as well as for improvements in quality and yield of crop plants and ornamentals. Thus, the invention contributes to the production of high quality and high yield agricultural products; for example, fruits, ornamentals, vegetables, cereals, and field crops. Genetically-improved seeds and other plant products that are produced using plants expressing the genes and methods described herein also render farming possible in areas previously unsuitable for agricultural production. The mechanisms disclosed herein for increasing plant yield and productivity is expected to be ubiquitous throughout the plant kingdom.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

Brief Description of the Drawings

Figure 1A is a bar graph showing the specificity of plant hormone responses in the *Arabidopsis* mesophyll protoplast transient expression system.

5 **Figure 1B** is a graph depicting a cytokinin dose response for the *ARR6* promoter induction.

Figure 1C is a bar graph showing the induction of the *ARR6* promoter by different cytokinins.

10 **Figure 1D** shows that CKI1 activation of *ARR6-LUC* requires HK activity and phosphorelay. The photograph (top) shows an autoradiogram depicting the expression of CKI1 proteins. The bar graph (bottom) shows the results of transfected protoplasts that were incubated for 3 hours to allow effector gene expression before cytokinin treatment for another 3 hours.

15 **Figure 1E** shows that CRE1 confers cytokinin hypersensitivity on the activation of the *ARR6-LUC* reporter. The photograph (top) shows an autoradiogram depicting the expression of *Arabidopsis* HK proteins. The results of this incubation with cytokinin are shown in the bar graph (bottom).

Figure 1F shows a series of photomicrographs showing that CKI1 is localized at the plasma membrane.

20 **Figure 2A** demonstrates that AHP acts as a shuttle between the cytosol and nucleus in cytokinin signalling. The photograph (top) depicts an autoradiogram demonstrating expression of AHP proteins. The bar graph (bottom) shows the effect of cytokinin treatment on plasmid expression.

25 **Figure 2B** shows a series of photomicrographs demonstrating that cytokinin induces transient AHP translocation.

Figure 3A shows negative and positive regulation by ARRs. The photograph depicts an autoradiogram (top) showing the expression of *Arabidopsis* ARR proteins. The bar graph (bottom) shows the activity of *ARR6* promoter activity when cells were then incubated with or without 100 nM *t*-zeatin
30 for another 3 hours before the activity was measured.

Figure 3B shows that ARR2s act downstream of CKI1 in cytokinin signalling.

Figure 3C shows a series of photomicrographs demonstrating that *Arabidopsis* ARR2s are localized in the nucleus.

5 Figure 4A is a photograph showing that ectopic expression of ARR2 is sufficient to promote cytokinin responses in transgenic tissues and plants when exposed to auxin.

10 Figure 4B is a photograph showing that ectopic expression of ARR2 is sufficient to promote cytokinin responses in transgenic tissues and plants when exposed to both auxin and cytokinin.

Figure 4C is a photograph demonstrating that cytokinin signaling delays dark-induced senescence in transgenic *Arabidopsis* plants.

Figure 5 is a schematic of a model of the cytokinin signal transduction pathway in *Arabidopsis*.

15

Detailed Description

Overview

Cytokinins are essential plant hormones involved in shoot meristem and leaf formation, cell division, chloroplast biogenesis, and senescence. Although
20 hybrid histidine protein kinases (HKs) have been implicated in cytokinin perception in *Arabidopsis*, the action of HK receptors and downstream signalling pathways remain elusive in plant cells. To study the action of HK receptors and downstream signaling pathways, a mesophyll protoplast transient assay based on the transcription activation of a cytokinin primary response gene *ARR6* in
25 *Arabidopsis* was developed. The use of this high-throughput cell assay for functional genomic analysis of the two-component regulators has provided a powerful tool to decipher the first complete cytokinin signalling pathway. Rather than following the established eukaryotic HK and MAPK cascade paradigm,

plant two-component signalling can integrate multiple HK activities to common AHPs as cytoplasm and nuclear shuttles, and distinct nuclear ARR_s as master regulatory outputs.

As is described in more detail below, a new eukaryotic two-component signalling circuit that initiates cytokinin signalling by employing distinct hybrid HK activities at the plasma membrane has been identified. In particular, AHPs are shown to act as novel eukaryotic signalling shuttles between the cytoplasm and nucleus in a transient and cytokinin-dependent manner. The short signalling circuit reaches the nuclear target genes by de-repressing nuclear response regulators ARR₁, ARR₂, and ARR₁₀ as transcription activators. The cytokinin-inducible ARR₄, ARR₅, ARR₆, and ARR₇ act as transcription repressors, suggesting a negative feedback loop. Thus, the *Arabidopsis* cytokinin signal transduction pathway consists of four major steps: HK sensing and signalling, AHP translocation, ARR de-repression, and a negative feedback loop by cytokinin-inducible ARR_s. Transgenic tissue and plant analyses support the importance of this core signalling pathway in diverse cytokinin responses. Furthermore, ectopic expression of the master regulator ARR₂ as the rate-limiting two-component output in transgenic *Arabidopsis* was sufficient to mimic cytokinin in promoting shoot meristem proliferation, leaf differentiation, expansion, and longevity.

The examples provided below are for the purposes of illustrating the invention, and should not be construed as limiting.

Cytokinin Signal is Perceived and Initiated by the Multiple HK receptors

To elucidate the regulatory circuitry in cytokinin signal transduction, a leaf cell assay based on cytokinin inducible transcription in *Arabidopsis* mesophyll protoplasts was developed. The 2.4 kb promoter of an *Arabidopsis* cytokinin primary response gene (SEQ ID NO.: 1), encoding the response regulator 6 (ARR₆), was fused to the firefly luciferase (LUC) reporter. In Fig. 1A, transfected protoplasts were incubated with or without 100 nM *t*-zeatin, 1 μ M

indole-3-acetic acid (IAA), or 100 μ M ABA. After 3 hours of incubation, LUC and GUS activities were determined. In transfected protoplasts, the activity of *ARR6-LUC* was specifically induced by cytokinin but not by other plant hormones such as abscisic acid (ABA) or auxin (Fig. 1A).

5 In the same system, the *GH3* promoter was only activated by auxin, whereas the *RD29A* promoter was solely induced by abscisic acid (ABA), demonstrating the specificity of three plant hormone responses in *Arabidopsis* mesophyll protoplasts (Fig. 1A). The activity of *ARR6-LUC* was induced by physiological concentration of a natural cytokinin *trans*-zeatin (*t*-zeatin) from 1 to
10 100 nM (Fig. 1B).

To further show the specificity of the cytokinin response, various active and inactive cytokinin analogues were examined. Only the active cytokinins, *t*-zeatin, 2-isopentenyladenine (2-IP; 100 nM) and benzyladenine (BA; 100 nM), induced the reporter gene *ARR6-LUC* (Fig. 1C).

15 The response in the protoplast system is similar to the cytokinin activation of various *ARR* genes shown *in planta* (D'Agostino et al., *supra*; Kiba et al., *supra*). Thus, we have established a reliable and physiological system to dissect the regulatory components in the cytokinin signal transduction pathway in *Arabidopsis*.

20

Cytokinin Signalling by Distinct HK Receptors and Phosphorelay

Despite the genetic identification of histidine kinases (HKs) as receptors for the plant hormones ethylene and cytokinin in *Arabidopsis*, whether or where HK activity is important for signal transduction in plant cells remains an
25 unresolved question due to the lack of physiological plant cell assays. CKI1, a hybrid HK with a conserved response regulator (RR) receiver domain has been implicated in cytokinin responses (Kakimoto, *supra*). The enhanced CKI1 expression in *Arabidopsis shooty callus* mutants also supports its role in cytokinin signalling (Frank et al., *Plant Physiol.* 122: 721-9, 2000). However, the
30 biochemical mechanism of CKI1 function and its role in cytokinin signalling are

still unknown. To determine whether CKI1 mediates cytokinin responsive transcription, the full-length *CKI1* gene (SEQ ID NO.: 13) was tagged with double haemagglutinin epitope (DHA) sequence and fused to a constitutive promoter, and co-transfected with *ARR6-LUC* into the *Arabidopsis* protoplasts.

5 Specifically, protoplasts were cotransfected with the *ARR6-LUC* reporter and the effector plasmid expressing CKI1 (SEQ ID NO.: 13), CKI1(H405Q) (SEQ ID NO.: 14), CKI1(D1050N) (SEQ ID NO.: 15), CKI1-GFP, or mER7-GFP (control). As shown in Fig. 1D, the wild-type CKI1 activated the *ARR6* promoter without exogenous *t*-zeatin. It is possible that over-expression of CKI1 renders
10 protoplasts hypersensitive to endogenous cytokinin and/or exceeds the capacity of negative regulators. Alternatively, CKI1 could encode a constitutively active HK connected to the cytokinin signal transduction pathway. Cytokinin treatment hardly enhanced the reporter gene activity in the presence of CKI1 (Fig. 1D).

To determine whether the HK activity and phosphorelay are required for
15 CKI1 activation of the *ARR6* promoter, the conserved His 405 and Asp 1040 residues in the HK and phosphorylation sites, respectively, were mutated. Specifically, protoplasts were transfected with plasmid DNA expressing CKI1 (SEQ ID NO.: 13), CKI (H405Q) (SEQ ID NO.: 14) and CKI1(D1040N) (SEQ ID NO.: 15), or vector DNA as a control. The [³⁵S]methionine labeled protein
20 from transfected protoplasts was immunoprecipitated with anti-HA and protein A-agarose, and analyzed by SDS-PAGE. Despite comparable expression of the CKI1(H405Q) and CKI1(D1050N) mutants as the wild-type CKI1, neither could activate the expression of *ARR6-LUC* (Fig. 1D). Furthermore, the CKI1(H405Q) mutant served as a strong dominant negative mutant to block the activation of the
25 *ARR6* promoter by exogenous *t*-zeatin. The CKI1(D1040N) mutant is less potent. These results indicated that HK activity and phosphorelay from His 405 to D1040 are required for the CKI1 function in activating cytokinin signalling. Moreover, the dominant negative CKI1(H405Q) mutant might interfere with cytokinin perception and/or disturb downstream signalling.

Recently, another *Arabidopsis* hybrid HK, CRE1/AHK4/WOL, has been shown to be a cytokinin receptor. The evidence is based on the inability of the *cre1* mutant to respond to cytokinin in the shoot induction assay and its ability to complement HK mutants of budding and fission yeast and *E. coli* in a cytokinin-dependent manner (Suzuki et al., *supra*; Inoue et al., *supra*; Mahonen et al., *Genes Dev.* 14: 2938-43, 2000). Since CRE1/AHK4/WOL is predominantly expressed in roots (Mahonen et al., *supra*; Ueguchi et al., *Plant Cell Physiol.* 42: 231-5, 2001) and the *cre1* and *wol* recessive mutants lack an obvious leaf phenotype (Inoue et al., *supra*; Mahonen et al., *supra*; Ueguchi et al., *supra*), it is possible that other closely related *Arabidopsis* HK genes, such as AHK2 and AHK3, and CKI1 provide cytokinin receptor functions in promoting shoot meristem and leaf formation, and delaying leaf senescence (Inoue et al., *supra*; Ueguchi et al., *supra*).

To directly test the function of these HKs in cytokinin signalling, constructs expressing AHK2 (SEQ ID NO.: 16), AHK3 (SEQ ID NO.: 17), or CRE1(AHK4/WOL) (SEQ ID NO.: 18) were cotransfected into *Arabidopsis* protoplast with the cytokinin-inducible reporter *ARR6-LUC*. Specifically, protoplasts were cotransfected with the *ARR6-LUC* reporter and the effector plasmid expressing AHK2, AHK3, CRE1(AHK4/WOL), CRE1(H467Q) (SEQ ID NO.: 19), CRE1(D973N) (SEQ ID NO.: 20), or mER7-GFP (control), and were analyzed after 6 hours of cytokinin treatment. Different from the CKI1 activation, over-expression of AHK2, AHK3, or CRE1(AHK4/WOL) did not enhance the expression of *ARR6-LUC* from the basal level in the absence of exogenous cytokinin. However, cytokinin treatment resulted in further activation of the reporter, especially for CRE1(AHK4/WOL) (Fig.1E). The requirement of the conserved His 467 and Asp 973 residues in the HK and the receiver domain, respectively, for the CRE1 activity was also tested. In Fig. 1E top, the expression of *Arabidopsis* HK proteins was detected by [³⁵S]methionine-labeled AHK2 (SEQ ID NO.: 26), AHK3 (SEQ ID NO.: 27), CRE1(AHK4/WOL) (SEQ ID NO.: 28), CRE1(H467Q) (SEQ ID NO.: 29), and CRE1(D973N) (SEQ ID NO.:

30) that were immunoprecipitated and analyzed by SDS-PAGE. CRE1(H467Q) and CRE1(D973N) mutants lost their ability to further enhance *ARR6-LUC* expression in the presence of exogenous cytokinin (Fig. 1E).

Moreover, both mutants also imposed a dominant negative effect on blocking cytokinin signalling. Similar to the CKI1 mutants, the His mutant is more potent than the Asp mutant, perhaps due to the flexible phosphorelay process between mutant and wild-type HKs. These results indicated that cytokinin signals are sensed by multiple HK receptors with different or overlapping expression patterns. Although CRE1 is predominantly expressed in roots (Mahonen et al., *supra*; Ueguchi et al., *supra*), it can certainly function in leaf cells as a cytokinin receptor. However, CKI1 and CRE1 represent two different types of cytokinin receptors that require HK activity and phosphorelay to initiate cytokinin signalling, but have different regulatory mechanisms and probably distinct affinity or specificity for cytokinins.

To gain insight into where the cytokinin signalling is perceived, the sub-cellular localization of CKI1 was examined using the CKI1-GFP fusion. We first confirmed that CKI1-GFP acted similarly as CKI1-HA in the protoplast assay based on *ARR6-LUC* activation (Fig. 1D). Confocal microscopy showed that CKI1 is mainly localized to the plasma membrane but not in the cytosol, ER or nucleus, indicated by various GFP markers (Fig. 1F). Specifically, protoplasts were transfected with CKI1-GFP or various GFP marker plasmid DNA. Tunicamycin (1 µg/ml) treatment was for 12 hours. The transfected protoplasts were analyzed with a confocal microscope. The subcellular GFP markers are “CAT-GFP” for the cytosol, “ER-GFP” for the endoplasmic reticulum, and “N-GFP” (HSP-GFP) for the nucleus. Since the N-terminus of CKI1 is replete with putative glycosylation sites, the effect of a glycosylation inhibitor, tunicamycin, in CKI1-GFP expression, localization or function was examined. Although tunicamycin did not affect the expression of other GFP constructs, the expression

of CKI1-GFP was completely abolished (Fig. 1F). This result showed that the glycosylation of CKI1 is important for its stability, processing and/or trafficking in plant cells.

5 HPs Acts as a Shuttle Between the Cytosol and Nucleus Shuttles in Cytokinin Signalling

The studies of multistep phosphorelay in yeast osmosensing and in various prokaryotic systems have shown the use of histidine-containing phosphotransmitters (HP) as signal transducers between receptors and response
10 regulators (Stock et al., *supra*; Posas et al., *supra*; Uhl and Miller, *EMBO J.* 15: 1028-36, 1996). The analysis of the *Arabidopsis* genome has revealed at least five genes encoding putative AHPs with unknown physiological functions. To determine whether AHPs are involved in the cytokinin signal transduction, AHP1, AHP2, and AHP5 were cloned into the plant expression vector and tested
15 in the protoplast cytokinin response assay based on the *ARR6-LUC* reporter activity. AHP1, AHP2, and AHP5 slightly increased the expression of the *ARR6-LUC* with or without *t*-zeatin (Fig. 2A). Specifically, protoplasts were cotransfected with the *ARR6-LUC* reporter and an effector plasmid carrying AHP1 (SEQ ID NO.: 21), AHP2 (SEQ ID NO.: 22), AHP5 (SEQ ID NO.: 23), or
20 AHP1 mutants (SEQ ID NOS.: 24 and 25). Transfected protoplasts were incubated for 6 hours to allow effector expression before cytokinin treatment for 3 hours. The AHP1 mutant, where the conserved His79 was replaced with Gln, did not repress the cytokinin activation of *ARR6-LUC*. Even the AHP1(H77Q, H79Q; SEQ ID NO.: 25) double mutant was not able to inhibit the *ARR6-LUC*
25 activation by cytokinin. AHPs and AHP1 mutants were expressed at comparable levels in the protoplast system (Fig. 2A top). This result indicated that AHPs are not the limiting factor or the key switch/regulators in cytokinin signalling. Their action as potential mediators may require cytokinin-dependent modification.

Previous *in vitro* studies have suggested that AHP1 and AHP2 could interact with the receiver domain of CKI1 (Nakamura et al., *Biosci. Biotechnol. Biochem.* 63: 1627-30, 1999).

To gain further *in vivo* support for the involvement of AHPs in cytokinin signalling, the action of AHP1-GFP in transfected protoplast in the absence or presence of cytokinin was followed using fluorescence microscopy. Specifically, protoplasts were transfected with *AHP1-GFP*, *AHP2-GFP*, or *AHP5-GFP* and incubated for 3 hours to allow sufficient expression. After *t*-zeatin (100 nM) treatment for 30 minutes, transfected cells were observed under a fluorescence microscope. As shown in Fig. 2B, AHP1-GFP was mainly localized in the cytoplasm without cytokinin stimulation, but was translocated into the nucleus by *t*-zeatin treatment. The cytokinin-dependent translocation of AHP1-GFP was transient, occurring within 30 to 90 minutes after cytokinin treatment (Fig. 2B). AHP2-GFP showed similar cytokinin-dependent translocation but not AHP5-GFP, suggesting their different functions in plant cells. Thus, AHP1 and AHP2 are believed to serve as shuttles and phosphorelay carriers between the cytokinin receptors and the downstream nuclear responses.

Opposite Regulation of Early Cytokinin Response Gene Transcription by Two Types of Nuclear RRs in Cytokinin Signalling

The requirement of HK activity and phosphorelay in the initiation of cytokinin signalling from CKI1 and CRE1, the cytokinin-inducible *ARR6* expression, the cytokinin-dependent translocation of AHPs into the nucleus, and the physical interaction of AHPs and ARRs in yeast (Suzuki et al., *Plant Cell Physiol.* 42: 37-45, 2001) suggest that some ARRs may act downstream in cytokinin signal transduction. There are two major subfamilies of *RR* genes in the *Arabidopsis* genome: the cytokinin-inducible A type and the DNA-binding B type (Urao, et al., *supra*; D'Agostino and Kieber, *Trends Biochem. Sci.* 24: 452-6, 1999; Sakai et al., *Plant J.* 24: 703-11, 2000). To investigate whether any *Arabidopsis* RRs are involved in cytokinin signalling, four representative A-type

ARR genes and three B-type *ARR* genes were cloned into a standard plant expression vector with an HA or GFP tag. Specifically, protoplasts were cotransfected with the *ARR6-LUC* reporter and the effector plasmid carrying *ARR1*, *ARR2*, *ARR4*, *ARR5*, *ARR6*, *ARR7*, *ARR10*, or various mutants, *ARR2* (D80N) (SEQ ID NO.: 10), *ARR4*(D95N) (SEQ ID NO.: 11), and *ARR6*(D76N) (SEQ ID NO.: 12). As shown in Fig. 3A, A-type ARRs, such as *ARR4*, *ARR5*, *ARR6*, and *ARR7*, repressed *ARR6-LUC* activity induced by 100 nM *t*-zeatin. Although *ARR6* seemed to be expressed at lower abundance, its repression activity was the strongest.

10 In contrast, the B-type ARRs, such as *ARR1*, *ARR2*, and *ARR10*, dramatically activated *ARR6-LUC* expression. Over-expression of *ARR1*, *ARR2*, and *ARR10* was sufficient to activate cytokinin signalling at different levels in the absence of exogenous cytokinin. *ARR1* and *ARR2* activated *ARR6-LUC* about 40- and 400-fold, respectively. Cytokinin treatment further enhanced the effect of *ARR2* on *ARR6-LUC* to over 1000-fold. The lower activation by *ARR1* could be due to its lower expression level in transfected protoplasts. In the absence of cytokinin, *ARR10* could activate *ARR6-LUC* about 10 fold. Cytokinin enhanced the effect of *ARR10* on the *ARR6-LUC* activity another 10-20 fold. The differential effect of *ARR2* and *ARR10* in activating cytokinin signalling could be attributed to differences in their intrinsic activities in DNA binding and/or transcription activation. It could also be due to their distinct affinity to endogenous repressors that likely exist in leaf protoplasts to prevent cytokinin signalling under unstimulated condition. Surprisingly, mutations of the conserved Asp residue in the receiver domains of *ARR4*(D95N), *ARR6*(D76N), and *ARR2*(D80N) did not lead to alteration of their repression and activation functions, respectively, in cytokinin signalling (Fig. 3A). The transfected cells were incubated for 3 hours to allow effector protein expression. This result is consistent with the model that phosphorelay stimulated by cytokinin may be involved in de-repression of the positive regulators like *ARR1*, *ARR2*, and

15
20
25

ARR10. Over-expression of these ARR2s likely bypasses the negative regulation and causes constitutive cytokinin responses without the signal.

To further support the role of ARR2 in cytokinin signalling, we designed a dominant mutant of ARR2 (Δ ARR2) (SEQ ID NO.: 4) containing only the DNA binding domain but not the putative transcription activation and the receiver domains (Sakai et al., *supra*). If Δ ARR2 can compete with the endogenous ARR2s in binding to the *ARR6* promoter without transcription activation, it may block cytokinin- or CKI1-dependent activation of *ARR6-LUC*. In Fig. 3B, protoplasts were cotransfected with the *ARR6-LUC* reporter alone or with the effector plasmids as indicated. Transfected protoplasts were incubated for 6 hours to allow effector protein expression before treatment with 100 nM *t*-zeatin for 3 hours. As shown in Fig. 3B, the dominant negative mutant of ARR2 could effectively block *ARR6-LUC* expression in the presence of 100 nM *t*-zeatin or CKI1, suggesting that ARR2 and/or ARR2-like proteins act downstream of cytokinin and CKI1-initiated phosphorelay as the nuclear targets in the cytokinin signal transduction pathway. Furthermore, the dominant negative effect of the CKI1(H450Q) mutant on the protoplast response to exogenous cytokinin could be bypassed by over-expression of the wild-type ARR2 (Fig. 3B). These epistasis analyses place ARR2 downstream of multiple cytokinin HK receptors. The finding of multiple ARR2 binding motifs, (G/A)GAT(T/C), in the promoter regions of *ARR6* and other cytokinin-inducible genes suggests that ARR2 could be a master regulator in cytokinin signalling (*Id.*). Exogenous cytokinin-initiated phosphorelay through CRE1, or AHK2 and AHK3 likely converges with the CKI1 phosphorelay on the common AHP shuttles and the nuclear ARR transcription activators.

To provide more evidence that the A-type and B-type ARR2s are transcription repressors and activators, respectively, in cytokinin signalling, the sub-cellular localization of ARR6-GFP and ARR2-GFP was examined. In Fig. 3C, Protoplasts were transfected with *ARR2-GFP*, *ARR2(D80N)-GFP*, Δ ARR2-GFP, or *ARR6-GFP* plasmid DNA and observed after 3-6 hours of incubation

with a fluorescence microscope. Both types of ARR2s are exclusively localized in the nucleus regardless of the cytokinin treatment (Fig. 3C). Their nuclear localization is likely independent of the phosphorylation state because ARR2(D80N)-GFP was found in the nucleus. The ARR2 dominant negative mutant (Δ ARR2) was also localized in the nucleus (Fig. 3C). Thus, the cytokinin-dependent phosphorelay does not play a role in the nuclear localization and DNA-binding of ARR2s or their intrinsic transcription activation or repression activities. The results support the model that cytokinin-initiated phosphorelay plays a role in releasing the sequestered activation-type ARR transcription factors from the yet unknown repressors (Fig. 5).

As depicted in Fig. 5, cytokinin signal is perceived internally or externally by multiple HKs at the plasma membrane (AHK2, AHK3, CKI1, or CRE1). Upon perception of the cytokinin signal, HKs initiate a signalling cascade via the phosphorelay that in turn results in the nuclear translocation of AHPs from the cytosol. Activated AHPs interact with sequestered ARR2s or ARR2 complex and release the activation type of ARR2s from putative repressor in the nucleus. The de-repressed ARR2s bind to multiple *cis* elements in the promoter of target genes. The transcription activation is essential for cell proliferation, shoot formation, and delayed senescence. The activation of the repressor-type of *ARR2s* as early cytokinin response genes provides a negative feedback mechanism that allows plant cells for fine-tuning or resetting the physiological state in diverse differentiation processes during plant growth and development (RD, response domain; BD, DNA binding domain; AD, transcription activation domain; PM, plasma membrane; N, nucleus; R, repressor).

This model is consistent with the findings that the N-terminus of ARR1 or ARR2 inhibits transcription based on a synthetic promoter (*Id.*), and the addition of cytokinin further enhances the effects of ARR1, ARR2 and ARR10 over-expression on the *ARR6-LUC* activity (Fig. 3). The activation of the repressor-

type of ARRs is thought to provide a negative feedback loop in controlling the transient induction of cytokinin early response genes and allows resetting and/or fine-tuning of the physiological state of the cells.

5 Ectopic expression of the master regulator ARR2 mimics cytokinin responses

In tissue culture, induction of cell proliferation and subsequent shoot formation require cytokinin. To determine whether the same cytokinin signalling pathway is responsible for transcription regulation, cell proliferation, shoot meristem initiation, and leaf formation, we developed a seedling cytokinin response assay. In Figs. 4A and 4B, *Arabidopsis* seedlings were transformed with *Agrobacteria* carrying *GFP*, *CKI1*, *CKI1(H405Q)*, *ARR2*, and *ARR6* constructs in the mini-binary vector pCB302 (Xiang et al., *Plant Mol. Biol.* 40: 711-7, 1999). Seedlings transformed with the *Agrobacteria* carrying the binary vector pCB301 without a Bar selection marker were used as a transformation control (vector).

15 The transformed seedlings were selected with glufosinate ammonium (10 µg/ml) and maintained on medium with auxin (IAA) only (Fig. 4A, +IAA, -2IP) or with auxin and cytokinin (Fig. 4B, +IAA, +2IP) for 14 days. The transformed tissues were mostly derived from the shoot apical meristem. The GFP control showed green callus formation with exogenous IAA but promoted shoot formation with

20 exogenous cytokinin. Both CKI1 and ARR2 promoted extensive cell proliferation and shoot formation even without exogenous cytokinin. The experiments were repeated three times with similar results. Compared with the GFP control, the CKI1(H405Q) mutant and ARR6 slightly inhibited cell proliferation on the same medium without or with exogenous cytokinin (Figs. 4A

25 and 4B). The results of this cytokinin seedling assay are consistent with those from protoplast transient expression analyses based on transcription. It is therefore clear that the HK activity of CKI1 is required to perceive and/or initiate cytokinin signalling.

To observe the consequences of constitutive cytokinin signalling *in planta*, the binary vectors expressing CKI1, CKI1(H405Q), ARR2, and ARR6 were

30

introduced into *Arabidopsis* plants. Ectopic expression of ARR2 in transgenic *Arabidopsis* plants displayed a spectrum of cytokinin-related phenotypes, including reduced apical dominance, increased leaf number and size, altered leaf shape and color, and delayed leaf senescence (Fig. 4C). Transgenic *Arabidopsis* plants carrying the CKI1 construct showed similar phenotypes, but not the CKI1(H405Q) and ARR6 transgenic plants serving as controls. Taken together, CKI1 and ARR2 are positive regulators in the cytokinin signal transduction pathway mediating transcription, cell proliferation, and shoot and leaf formation, as well as delaying leaf senescence. ARR2 could be one of the direct downstream targets of CKI1 and CRE1, and serve as a master regulator in diverse cytokinin responses.

Materials and Methods

The above-mentioned results were obtained using the following materials and methods.

Plasmid Constructs

The 2.4 kb *Arabidopsis* ARR6 promoter was fused to the firefly luciferase gene to create the ARR6-LUC reporter construct. The CKI1 and AHK2 genes were amplified from the *Arabidopsis* genomic DNA by polymerase chain reaction (PCR). The AHK3, CRE1(CRE1/WOL), AHP1, AHP2, AHP5 and ARR1, ARR 2, ARR4, ARR5, ARR6, ARR7, and ARR10 coding regions were obtained by PCR from an *Arabidopsis* cDNA library (Sheen, *Science* 274: 1900-2, 1996). All of the mutants were generated by QuickChange site-directed mutagenesis (Stratagene). The coding regions of all proteins were tagged with either two copies of the haemagglutinin epitope (DHA) or GFP and inserted into a plant expression vector containing the 35SC4PPDK promoter (Sheen, *supra*) and the nos terminator. All PCR products and the mutations were confirmed by DNA sequencing.

Arabidopsis Protoplast Transient Expression Assay

Arabidopsis Bensheim (BE) protoplasts were isolated and transfected as described with some modifications (Kovtun et al., *Proc. Natl. Acad. Sci. U S A* 97: 2940-5 2000). Typically, 2×10^4 protoplasts were transfected with 10 μ g of
5 plasmid carrying different combinations of a reporter, effectors, and an internal control. For instance, in the experiments shown in Fig. 1A, protoplasts (4×10^4) were transfected with 2 μ g of *UBQ10-GUS* (internal control) and 20 μ g of *ARR6-LUC*, *GH3-LUC*, or *RD29A-LUC* plasmid DNA. Transfected protoplasts were incubated at 1×10^4 per ml with/without 1-100 nM of *t*-zeatin for 3-6 hours under
10 the light at 23 °C. The *UBQ10-GUS* construct was used as an internal control to normalize the variations of each transfection in cell numbers, transformation efficiency and cell viability. The results are shown as the means of duplicate samples with the standard deviation. All transient experiments were repeated at least three times with similar results. GFP fluorescence was observed by either
15 Nikon TE200 fluorescent microscopy or Leica TCSNT confocal microscopy.

Protein Immunoprecipitation

Transfected protoplasts were incubated with [35 S]-methione (200 μ Ci/ml) for 6 hours. The effector proteins were immunoprecipitated as described (Sheen,
20 *supra*), analyzed by SDS/PAGE (10 or 12.5 %) and visualized by fluorography.

Seedling Cytokinin Response Assay

Four-day-old *Arabidopsis* BE seedlings were placed on the callus induction medium, containing 0.5 mg/L 2,4-D, 0.05 mg/L BAP, and 0.05 mg/L
25 kinetin, for four days. The seedlings were co-cultivated and transformed with *Agrobacterium tumefaciens* GV3101 carrying pCB302 with either *CK11*, *CK11(H405Q)*, *ARR2*, *ARR6*, *GFP* under control of the 35SC4PPDK promoter, or the pCB301 binary vector (Xiang et al., *Plant Mol. Biol.* 40:711-7, 1999) alone for three days. Transformed seedlings were then selected on the glufosinate
30 ammonium-containing medium (with either 0.15 mg/L IAA alone or with 0.15

mg/L IAA and 5 mg/L 2IP) and observed up to 4 weeks. The transformed proliferating tissues were mostly derived from the apical shoot meristem of the seedlings.

5 Transgenic plant analysis

The same constructs analyzed in the *Arabidopsis* protoplast transient expression and seedling assays, including *CKI1*, *CKI1(H405Q)*, *ARR2*, and *ARR6*, were used to generate *Arabidopsis* transgenic plants using the floral dip method and Bar selection as described (Clough and Bent, *Plant J.* 16: 735-43, 1998).

- 10 More than 200 transgenic plants were obtained with each construct. The *ARR2* and *CKI1* transgenic plants developed branches early, produced more leaves that are bigger and greener but flowered later than the control *CKI1(H405Q)* and *ARR6* transgenic plants. For senescence assay, fully expanded fourth leaves from representative transgenic plants were detached and floated on distilled water in
15 the dark for four days.

Isolation of Sequences Encoding A-type and B-type Response Regulators or Histidine Kinases

- The isolation of additional genes encoding A-type and B-type response
20 regulators or histidine kinase genes is accomplished using standard strategies and techniques that are well known in the art.

- In one particular example, the B-type response regulator sequence (for example, *ARR1*, *ARR2*, or *ARR10* sequences), the A-type response regulator sequence, (for example, *ARR4*, *ARR5*, *ARR6*, or *ARR7*), or the histidine kinase
25 (for example, *CRE1* or *CKI1*), described herein may be used, together with conventional screening methods of nucleic acid hybridization screening, to isolate additional sequences encoding these regulators. Such hybridization techniques and screening procedures are well known to those skilled in the art and are described, for example, in Benton and Davis, *Science* 196: 180, 1977; Grunstein
30 and Hogness, *Proc. Natl. Acad. Sci., USA* 72: 3961, 1975; Ausubel et al. *Current*

Protocols in Molecular Biology, Wiley Interscience, New York; Berger and Kimmel, *Guide to Molecular Cloning Techniques*, 1987, Academic Press, New York.; and Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, New York. In one particular example, all or part
5 of the response regulator gene may be used as a probe to screen a recombinant plant DNA library for genes having sequence identity or similarity to the response regulator gene. Hybridizing sequences are detected by plaque or colony hybridization according to the methods described below.

Alternatively, using all or a portion of the amino acid sequence of a
10 response regulator, one may readily design oligonucleotide probes, including degenerate oligonucleotide probes (i.e., a mixture of all possible coding sequences for a given amino acid sequence). These oligonucleotides may be based upon the sequence of either DNA strand and any appropriate portion of the response regulator sequence. General methods for designing and preparing such
15 probes are provided, for example, in Ausubel et al., *supra*; and Berger and Kimmel, *supra*. These oligonucleotides are useful for response regulator or histidine kinase sequence isolation, either through their use as probes capable of hybridizing to response regulator complementary sequences or as primers for various amplification techniques, for example, polymerase chain reaction (PCR)
20 cloning strategies. If desired, a combination of different oligonucleotide probes may be used for the screening of a recombinant DNA library. The oligonucleotides may be detectably-labeled using methods known in the art and used to probe filter replicas from a recombinant DNA library. Recombinant DNA libraries are prepared according to methods well known in the art, for
25 example, as described in Ausubel et al. (*supra*), or they may be obtained from commercial sources.

As discussed above, response regulator-specific oligonucleotides may also be used as primers in amplification cloning strategies, for example, using PCR. PCR methods are well known in the art and are described, for example, in *PCR Technology*, Erlich, ed., Stockton Press, London, 1989; *PCR Protocols: A Guide*
30

to *Methods and Applications*, Innis et al., eds., Academic Press, Inc., New York, 1990; and Ausubel et al. (*supra*). Primers are optionally designed to allow cloning of the amplified product into a suitable vector, for example, by including appropriate restriction sites at the 5' and 3' ends of the amplified fragment (as
5 described herein). If desired, response regulator sequences may be isolated using the PCR "RACE" technique, or Rapid Amplification of cDNA Ends (see, e.g., Innis et al. (*supra*)). By this method, oligonucleotide primers based on the response regulator sequence are oriented in the 3' and 5' directions and are used to generate overlapping PCR fragments. These overlapping 3'- and 5'-end RACE
10 products are combined to produce an intact full-length cDNA. This method is described in Innis et al. (*supra*); and Frohman et al., *Proc. Natl. Acad. Sci. USA* 85: 8998, 1988.

Confirmation of a sequence's relatedness to the response regulators may be accomplished by a variety of conventional methods including, but not limited to,
15 sequence comparison of the gene and its expressed product to known response regulators for example those disclosed herein. In addition, the activity of the gene product may be evaluated according to any of the techniques described herein.

Once a response regulator gene is identified, it is cloned according to
20 standard methods and used for the construction of plant expression vectors as described below.

Expression Constructs

Those skilled in the field of molecular biology will understand that any
25 of a wide variety of expression systems may be used to express the response regulators disclosed herein. For example, an A-type or B-type response regulator may be expressed for the desired effect in any of a number of plant hosts including, without limitation, algae, tree species, ornamental species, temperate fruit species, tropical fruit species, vegetable species, legume species, crucifer
30 species, monocots, dicots, or in any plant of commercial or agricultural

significance. Particular examples of suitable plant hosts include, but are not limited to, Conifers, Petunia, Tomato, Potato, Tobacco, *Arabidopsis*, Lettuce, Sunflower, Oilseed rape, Flax, Cotton, Sugarbeet, Celery, Soybean, Alfalfa, *Medicago*, Lotus, *Vigna*, Cucumber, Carrot, Eggplant, Cauliflower, Horseradish, Morning Glory, Poplar, Walnut, Apple, Grape, Asparagus, Rice, Maize, Millet, Onion, Barley, Orchard grass, Oat, Rye, and Wheat. In addition, as is discussed below, transgenic expression constructs may be expressed in a plant to increase plants yield, shoot formation, leaf growth, photosynthesis, or delay senescence.

Materials for expressing these genes are available from a wide range of sources including the American Type Culture Collection (Rockland, MD); or from any of a number seed companies, for example, W. Atlee Burpee Seed Co. (Warminster, PA), Park Seed Co. (Greenwood, SC), Johnny Seed Co. (Albion, ME), or Northrup King Seeds (Harstville, SC). Descriptions and sources of useful host cells are also found in Vasil, Cell Culture and Somatic Cell Genetics of Plants, Vol I, II, III Laboratory Procedures and Their Applications Academic Press, New York, 1984; Dixon, Plant Cell Culture-A Practical Approach, IRL Press, Oxford University, 1985; Green et al., Plant Tissue and Cell Culture, Academic Press, New York, 1987; and Gasser and Fraley, Science 244: 1293, 1989.

The method of transformation or transfection and the choice of vehicle for expression of the A-type or B-type response regulators, or a histidine kinase will depend on the host system selected. Transformation and transfection methods are described, e.g., in Ausubel et al. (*supra*); Weissbach and Weissbach, Methods for Plant Molecular Biology, Academic Press, 1989; Gelvin et al., Plant Molecular Biology Manual, Kluwer Academic Publishers, 1990; Kindle, *Proc. Natl. Acad. Sci., USA* 87: 1228, 1990; Potrykus, *Annu. Rev. Plant Physiol. Plant Mol. Biology* 42: 205, 1991; and BioRad (Hercules, CA) Technical Bulletin #1687 (Biolistic Particle Delivery Systems). Expression vehicles may be chosen from those provided, e.g., in Cloning Vectors: A Laboratory Manual (Pouwels et al., 1985, Supp. 1987); Gasser and Fraley (*supra*); Clontech Molecular Biology

Catalog (Catalog 1992/93 Tools for the Molecular Biologist, Palo Alto, CA); and the references cited above. Other expression constructs are described by Fraley et al. (U.S. Pat. No. 5,352,605).

Most preferably, an A-type or B-type response regulator or a histidine kinase is produced by a stably-transfected plant cell line, a transiently-transfected plant cell line, or by a transgenic plant. A number of vectors suitable for stable transfection of plant cells or for the establishment of transgenic plants are available to the public; such vectors are described in Pouwels et al. (*supra*), Weissbach and Weissbach (*supra*), and Gelvin et al. (*supra*). Methods for constructing such cell lines are described in, e.g., Weissbach and Weissbach (*supra*), and Gelvin et al. (*supra*). Typically, plant expression vectors include (1) a cloned plant gene under the transcriptional control of 5' and 3' regulatory sequences and (2) a dominant selectable marker. Such plant expression vectors may also contain, if desired, a promoter regulatory region (for example, one conferring inducible or constitutive, pathogen- or wound-induced, environmentally- or developmentally-regulated, or cell- or tissue-specific expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

Once the desired nucleic acid sequence encoding an A-type or B-type response regulators or a histidine kinase is obtained as described above, it may be manipulated in a variety of ways known in the art. For example, where the sequence involves non-coding flanking regions, the flanking regions may be subjected to mutagenesis.

In general, the constructs will involve regulatory regions functional in plants which provide for modified production of the regulator protein or regulator transcript (such as an antisense transcript) as discussed herein. The open reading frame coding for the regulator protein or functional fragment thereof will be joined at its 5' end to a transcription initiation regulatory region such as the

30

sequence naturally found in the 5' upstream region of the A-type or B-type response regulator, or a histidine kinase. Numerous other transcription initiation regions are available which provide for constitutive or inducible regulation.

For applications where developmental, cell, tissue, hormonal, or
5 environmental expression is desired, appropriate 5' upstream non-coding regions are obtained from other genes, for example, from genes regulated during meristem development, seed development, embryo development, or leaf development.

Regulatory transcript termination regions may also be provided in
10 DNA constructs of this invention as well. Transcript termination regions may be provided by the DNA sequence encoding the A-type or B-type response regulator or histidine kinase, or any convenient transcription termination region derived from a different gene source. The transcript termination region will contain preferably at least 1-3 kb of sequence 3' to the structural gene from which the
15 termination region is derived. Plant expression constructs having, for example, A-type or B-type response regulator as the DNA sequence of interest for expression may be employed with a wide variety of plant life. Such genetically-engineered plants are useful for a variety of industrial and agricultural applications as discussed herein. Importantly, this invention is applicable to
20 dicotyledons and monocotyledons, and will be readily applicable to any new or improved transformation or regeneration method.

An example of a useful plant promoter according to the invention is a caulimovirus promoter, for example, a cauliflower mosaic virus (CaMV)
promoter. These promoters confer high levels of expression in most plant tissues,
25 and the activity of these promoters is not dependent on virally encoded proteins. CaMV is a source for both the 35S and 19S promoters. In most tissues of transgenic plants, the CaMV 35S promoter is a strong promoter (see, e.g., Odell et al., *Nature* 313: 810, 1985). The CaMV promoter is also highly active in monocots (see, e.g., Dekeyser et al., *Plant Cell* 2: 591, 1990; Terada and
30 Shimamoto, *Mol. Gen. Genet.* 220: 389, 1990). Moreover, activity of this

promoter can be further increased (i.e., between 2-10 fold) by duplication of the CaMV 35S promoter (see e.g., Kay et al., *Science* 236: 1299, 1987; Ow et al., *Proc. Natl. Acad. Sci., U.S.A.* 84: 4870, 1987; and Fang et al., *Plant Cell* 1: 141, 1989). In addition, the a minimal 35S promoter may also be used as is described
5 herein.

Other useful plant promoters include, without limitation, the nopaline synthase promoter (An et al., *Plant Physiol.* 88: 547, 1988) and the octopine synthase promoter (Fromm et al., *Plant Cell* 1: 977, 1989).

For certain applications, it may be desirable to produce the regulator in
10 an appropriate tissue, at an appropriate level, or at an appropriate developmental time. For this purpose, there are an assortment of gene promoters, each with its own distinct characteristics embodied in its regulatory sequences, shown to be regulated in response to the environment, hormones, and/or developmental cues. These include gene promoters that are responsible for heat-regulated gene
15 expression (see, e.g., Callis et al., *Plant Physiol.* 88: 965, 1988; Takahashi and Komeda, *Mol. Gen. Genet.* 219: 365, 1989; and Takahashi et al., *Plant J.* 2: 751, 1992), light-regulated gene expression (e.g., the *Arabidopsis Cab2* photosynthetic, leaf specific promoter described by Mitra et al., *Plant Mol. Biol.* 12: 169-179, 1989; the pea *rbcS-3A* described by Kuhlemeier et al., *Plant Cell* 1:
20 471, 1989; the maize *rbcS* promoter described by Schäffner and Sheen, *Plant Cell* 3: 997, 1991; or the chlorophyll a/b-binding protein gene found in pea described by Simpson et al., *EMBO J.* 4: 2723, 1985), hormone-regulated gene expression (for example, the abscisic acid (ABA) responsive sequences from the *Em* gene of wheat described by Marcotte et al., *Plant Cell* 1: 969, 1989; the ABA-inducible
25 HVA1 and HVA22, and rd29A promoters described for barley and *Arabidopsis* by Straub et al., *Plant Cell* 6: 617, 1994, Shen et al., *Plant Cell* 7: 295, 1995), wound-induced gene expression (for example, of *wun1* described by Siebertz et al., *Plant Cell* 1: 961, 1989), and organ-specific gene expression (for example, of the tuber-specific storage protein gene described by Roshal et al., *EMBO J.* 6:
30 1155, 1987; the 23-kDa zein gene from maize described by Scherthaner et al.,

EMBO J. 7: 1249, 1988; or the French bean β -phaseolin gene described by Bustos et al., *Plant Cell* 1: 839, 1989; the vegetative storage protein promoter (soybean vspB) described by Sadka et al. (*Plant Cell* 6: 737-749, 1994)), cycling promoters (e.g., the *Arabidopsis cdc2a* promoter described by Hemerly et al., *Proc. Natl. Acad. Sci. USA* 89: 3295-3299, 1992), senescence-specific promoters (e.g., the *Arabidopsis* SAG12 promoter described by Gan et al., *Science*: 270, 1986-1988, 1995), seed-specific promoters (for example, endosperm-specific or embryo-specific promoters), or pathogen-inducible promoters (for example, PR-1 or β -1,3 glucanase promoters).

10 Plant expression vectors may also optionally include RNA processing signals, e.g, introns, which have been shown to be important for efficient RNA synthesis and accumulation (Callis et al., *Genes and Dev.* 1: 1183, 1987). The location of the RNA splice sequences can dramatically influence the level of transgene expression in plants. In view of this fact, an intron may be positioned
15 upstream or downstream of the regulator encoding sequence in the transgene to modulate levels of gene expression.

In addition to the aforementioned 5' regulatory control sequences, the expression vectors may also include regulatory control regions which are generally present in the 3' regions of plant genes (Thornburg et al., *Proc. Natl. Acad. Sci. U.S.A.* 84: 744, 1987; An et al., *Plant Cell* 1: 115, 1989). For
20 example, the 3' terminator region may be included in the expression vector to increase stability of the mRNA. One such terminator region may be derived from the PI-II terminator region of potato. In addition, other commonly used terminators are derived from the octopine or nopaline synthase signals.

25 The plant expression vector also typically contains a dominant selectable marker gene used to identify those cells that have become transformed. Useful selectable genes for plant systems include genes encoding antibiotic resistance, for example, those encoding resistance to hygromycin, kanamycin, bleomycin, G418, streptomycin, or spectinomycin. Genes required for
30 photosynthesis may also be used as selectable markers in photosynthetic-deficient

strains. Finally, genes encoding herbicide resistance may be used as selectable markers; useful herbicide resistance genes include the *bar* gene encoding the enzyme phosphinothricin acetyltransferase and conferring resistance to the broad spectrum herbicide Basta® (Hoechst AG, Frankfurt, Germany).

5 Efficient use of selectable markers is facilitated by a determination of the susceptibility of a plant cell to a particular selectable agent and a determination of the concentration of this agent which effectively kills most, if not all, of the untransformed cells. Some useful concentrations of antibiotics for tobacco transformation include, e.g., 75-100 µg/mL (kanamycin), 20-50 µg/mL
10 (hygromycin), or 5-10 µg/mL (bleomycin). A useful strategy for selection of transformants for herbicide resistance is described, e.g., by Vasil et al., *supra*.

 It should be readily apparent to one skilled in the art of molecular biology, especially in the field of plant molecular biology, that the level of gene expression is dependent, not only on the combination of promoters, RNA
15 processing signals, and terminator elements, but also on how these elements are used to increase the levels of selectable marker gene expression.

Plant Transformation

 Upon construction of the plant expression vector, several standard
20 methods are available for introduction of the vector into a plant host, thereby generating a transgenic plant. These methods include (1) *Agrobacterium*-mediated transformation (*A. tumefaciens* or *A. rhizogenes*) (see, e.g., Lichtenstein and Fuller, In: *Genetic Engineering*, Vol 6, Rigby, ed., London, Academic Press, 1987; and Lichtenstein and Draper, In: *DNA Cloning*, Vol II, Glover, ed., Oxford,
25 IRI Press, 1985), (2) the particle delivery system (see, e.g., Gordon-Kamm et al., *Plant Cell* 2: 603,1990); or BioRad Technical Bulletin #1687, *supra*), (3) microinjection protocols (see, e.g., Green et al., *supra*), (4) polyethylene glycol (PEG) procedures (see, e.g., Draper et al., *Plant Cell Physiol.* 23: 451, 1982; or e.g., Zhang and Wu, *Theor. Appl. Genet.* 76: 835, 1988), (5) liposome-mediated
30 DNA uptake (see, .g., Freeman et al., *Plant Cell Physiol.* 25: 1353, 1984), (6)

electroporation protocols (see, e.g., Gelvin et al., *supra*; Dekeyser et al., *supra*; Fromm et al., *Nature* 319: 791, 1986; Sheen, *Plant Cell* 2: 1027, 1990; or Jang and Sheen, *Plant Cell* 6: 1665, 1994), (7) the vortexing method (see, e.g., Kindle, *supra*), and floral dip method (see, e.g., Clough and Bent, *Plant J.* 16, 735-43.

5 (1998)). The method of transformation is not critical to the invention. Any method which provides for efficient transformation may be employed. As newer methods are available to transform crops or other host cells, they may be directly applied.

The following is an example outlining one particular technique, an
10 *Agrobacterium*-mediated plant transformation. By this technique, the general process for manipulating genes to be transferred into the genome of plant cells is carried out in two phases. First, cloning and DNA modification steps are carried out in *E. coli*, and the plasmid containing the gene construct of interest is transferred by conjugation or electroporation into *Agrobacterium*. Second, the
15 resulting *Agrobacterium* strain is used to transform plant cells. Thus, for the generalized plant expression vector, the plasmid contains an origin of replication that allows it to replicate in *Agrobacterium* and a high copy number origin of replication functional in *E. coli*. This permits facile production and testing of transgenes in *E. coli* prior to transfer to *Agrobacterium* for subsequent
20 introduction into plants. Resistance genes can be carried on the vector, one for selection in bacteria, for example, streptomycin, and another that will function in plants, for example, a gene encoding kanamycin resistance or herbicide resistance. Also present on the vector are restriction endonuclease sites for the addition of one or more transgenes and directional T-DNA border sequences
25 which, when recognized by the transfer functions of *Agrobacterium*, delimit the DNA region that will be transferred to the plant.

In another example, plant cells may be transformed by shooting into the cell tungsten microprojectiles on which cloned DNA is precipitated. In the Biolistic Apparatus (Bio-Rad) used for the shooting, a gunpowder charge (22
30 caliber Power Piston Tool Charge) or an air-driven blast drives a plastic

macroprojectile through a gun barrel. An aliquot of a suspension of tungsten particles on which DNA has been precipitated is placed on the front of the plastic macroprojectile. The latter is fired at an acrylic stopping plate that has a hole through it that is too small for the macroprojectile to pass through. As a result, the plastic macroprojectile smashes against the stopping plate, and the tungsten microprojectiles continue toward their target through the hole in the plate. For the instant invention the target can be any plant cell, tissue, seed, or embryo. The DNA introduced into the cell on the microprojectiles becomes integrated into either the nucleus or the chloroplast.

In general, transfer and expression of transgenes in plant cells are now routine practices to those skilled in the art, and have become major tools to carry out gene expression studies in plants and to produce improved plant varieties of agricultural or commercial interest.

Transgenic Plant Regeneration

Whole plants can be regenerated, for example, from single cells, callus tissue, or leaf discs transformed with a plant expression vector according to standard plant tissue culture techniques. It is well known in the art that various cells, tissues, and organs from almost any plant can be successfully cultured to regenerate an entire plant; such techniques are described, e.g., in Vasil, *supra*; Green et al., *supra*; Weissbach and Weissbach, *supra*; and Gelvin et al., *supra*.

In one particular example, a cloned B-type response regulator or histidine kinase construct under the control of the *nos* promoter and the nopaline synthase terminator and carrying a selectable marker (for example, kanamycin resistance) is transformed into *Agrobacterium*. Transformation of leaf discs (for example, of tobacco or potato leaf discs), with vector-containing *Agrobacterium* is carried out as described by Horsch et al. (*Science* 227: 1229, 1985). Putative transformants are selected after a few weeks (for example, 3 to 5 weeks) on plant tissue culture media containing kanamycin (e.g. 100 µg/mL). Kanamycin-resistant shoots are then placed on plant tissue culture media without hormones

for root initiation. Kanamycin-resistant plants are then selected for greenhouse growth. If desired, seeds from self-fertilized transgenic plants can then be sowed in a soil-less medium and grown in a greenhouse. Kanamycin-resistant progeny are selected by sowing surfaced sterilized seeds on hormone-free kanamycin-
5 containing media. Analysis for the integration of the transgene is accomplished by standard techniques (see, for example, Ausubel et al., *supra*; Gelvin et al., *supra*).

Transgenic plants expressing the selectable marker are then screened for transmission of the transgene DNA by standard immunoblot and DNA
10 detection techniques. Each positive transgenic plant and its transgenic progeny are unique in comparison to other transgenic plants established with the same transgene. Integration of the transgene DNA into the plant genomic DNA is in most cases random, and the site of integration can profoundly affect the levels and the tissue and developmental patterns of transgene expression. Consequently,
15 a number of transgenic lines are usually screened for each transgene to identify and select plants with the most appropriate expression profiles.

Transgenic lines are evaluated for levels of transgene expression. Expression at the RNA level is determined initially to identify and quantitate expression-positive plants. Standard techniques for RNA analysis are employed
20 and include PCR amplification assays using oligonucleotide primers designed to amplify only transgene RNA templates and solution hybridization assays using transgene-specific probes (see, e.g., Ausubel et al., *supra*). The RNA-positive plants are then analyzed for protein expression by Western immunoblot analysis using specific antibodies (see, e.g., Ausubel et al., *supra*). In addition, *in situ*
25 hybridization and immunocytochemistry according to standard protocols can be done using transgene-specific nucleotide probes and antibodies, respectively, to localize sites of expression within transgenic tissue.

Increasing Plant Yield

To test the ability of the genes described herein to improve crop yield or productivity, seeds of transgenic plants expressing a B-type response regulator or histidine kinase are planted in test plots, and their agronomic performance is compared to standard plants using techniques familiar to those of skill in the art. Optionally included in this comparison are plants of similar genetic background without the transgene. A yield benefit is observed and plants exhibiting the increased yield are advanced for commercialization.

In addition, transgenic plants expressing a B-type response regulator or histidine kinase are field tested for agronomic performance under conditions, including, but not limited to, increased leaf production. When compared to nontransgenic plants, transgenic plants expressing the B-type response regulator exhibit higher yield than their non-transgenic counterparts.

Increasing Plant Shoot Formation

To test the ability of the genes described herein to increase shoot formation, seedlings are transformed with a transgene expressing a B-type response regulator or histidine kinase and shoot formation is measured, for example, by evaluating the development of green callus formation, using techniques familiar to those of skill in the art. Optionally included in this comparison are plants of similar genetic background without the transgene. Plants exhibiting increased shoot formation relative to control plants are useful in the invention.

Delaying Plant Senescence

To test the ability of the genes described herein to delay senescence, fully expanded fourth leaves of plants expressing a recombinant B-type response regulator or histidine kinase are detached and floated on distilled water in the dark for four days, and their leaf size and quality of green color are compared

using techniques familiar to those of skill in the art. Optionally included in this comparison are plants of similar genetic background without the transgene.

Enhancing Plant Photosynthesis

5 To test the ability of the genes and domains described herein to enhance photosynthesis, seedlings are transformed with a transgene expressing a recombinant B-type response regulator or histidine kinase and photosynthesis measured using techniques familiar to those of skill in the art. Optionally included in this comparison are plants of similar genetic background without the
10 transgene.

Increasing Plant Leaf Growth

 To test the ability of the genes described herein to increase leaf growth, seedlings are transformed with a transgene expressing a recombinant B-type
15 response regulator and leaf growth measured, for example, by measuring leaf number and leaf area using techniques familiar to those of skill in the art. Optionally included in this comparison are plants of similar genetic background without the transgene.

20 Silencing A-type Response Regulator Gene Expression

 Plants having decreased expression of an A-type response regulator are useful, for example, for increasing plant productivity, delaying senescence, increasing photosynthesis, increased leaf growth, and increased productivity. Plants having decreased expression of such A-type response regulators are
25 generated according to standard gene silencing methods including, without limitation, co-suppression and antisense methodologies, expression of dominant negative gene products, and creation of plants having mutated genes encoding such regulators.

Co-Suppression

One preferred method of silencing gene expression is co-suppression (also referred to as sense suppression). This technique, which involves introduction of a nucleic acid configured in the sense orientation, has been shown to effectively
5 block the transcription of target genes (see for example, Napoli et al., *Plant Cell*, 2:279-289, 1990 and Jorgensen et al., U.S. Patent No. 5,034,323).

Generally, sense suppression involves transcription of the introduced sequence such as a gene encoding an A-type response regulator. However, co-suppression may also occur where the introduced sequence contains no coding
10 sequence per se, but only intron or untranslated sequences homologous to sequences present in the primary transcript of the endogenous gene to be repressed. The introduced sequence generally will be substantially identical to the endogenous gene targeted for repression. Such identity is typically greater than about 50%, but higher identities (for example, 80% or even 95%) are
15 preferred because they result in more effective repression. The effect of co-suppression may also be applied to other proteins within a similar family of genes exhibiting homology or substantial homology. Segments from a gene from one plant can be used directly, for example, to inhibit expression of homologous genes in different plant species.

20 In sense suppression, the introduced sequence, requiring less than absolute identity, need not be full length, relative to either the primary transcription product or to fully processed mRNA. A higher identity in a shorter than full length sequence compensates for a longer sequence with less identity.

Furthermore, the introduced sequence need not have the same intron or exon
25 pattern, and identity of non-coding segments may be equally effective. Sequences of at least 50 base pairs are preferred, with introduced sequences of greater length being more preferred (see, for example, those methods described by Jorgensen et al., *supra*).

Antisense Suppression

In antisense technology, a nucleic acid segment from the desired plant gene is cloned and operably linked to an expression control region such that the antisense strand of RNA is synthesized. The construct is then transformed into
5 plants and the antisense strand of RNA is produced. In plant cells, it has been shown that antisense RNA inhibits gene expression.

The nucleic acid segment to be introduced in antisense suppression is generally substantially identical to at least a portion of the endogenous gene or genes to be repressed, but need not be identical. The vectors of the present
10 invention therefore can be designed such that the inhibitory effect applies to other proteins within a family of genes exhibiting homology or substantial homology to the target gene. Segments from a gene from one plant can be used, for example, directly to inhibit expression of homologous genes in different plant species.

The introduced sequence also need not be full length relative to either the
15 primary transcription product or to fully processed mRNA. Generally, higher homology can be used to compensate for the use of a shorter sequence. Moreover, the introduced sequence need not have the same intron or exon pattern, and homology of non-coding segments will be equally effective. In general, such an antisense sequence will usually be at least 15 base pairs, preferably about 15-
20 200 base pairs, and more preferably 200-2,000 base pairs in length or greater. The antisense sequence may be complementary to all or a portion of the gene to be suppressed and as appreciated by those skilled in the art, the particular site or sites to which the antisense sequence binds as well as the length of the antisense sequence will vary, depending upon the degree of inhibition desired and the
25 uniqueness of the antisense sequence. A transcriptional construct expressing a plant A-type response regulator (e.g., those described herein) antisense nucleotide sequence includes, in the direction of transcription, a promoter, the sequence coding for the antisense RNA on the sense strand, and a transcriptional termination region. Antisense sequences may be constructed and expressed as
30 described, for example, in van der Krol et al. (*Gene* 72: 45-50, 1988); Rodermel

et al. (*Cell* 55: 673-681, 1988); Mol et al. (*FEBS Lett.* 268: 427-430, 1990); Weigel and Nilsson (*Nature* 377: 495-500, 1995); Cheung et al., (*Cell* 82: 383-393, 1995); and Shewmaker et al. (U.S. Patent No. 5,107,065).

5 In one working example, antisense expression of the gene encoding A-type response regulators of *Arabidopsis* or an A-type response regulator homolog is used to increase plant yield. In one particular approach, a plant expression vector is constructed that contains the cDNA sequence of A-type response regulator homolog in antisense orientation that is constitutively expressed under the control of the rice actin promoter described by Wu et al. (WO 91/09948). This
10 expression vector is then used to transform rice plants according to conventional methods, for example, using the methods described in Hiei et al. (*Plant J.* 6:271-282, 1994).

Dominant Negatives

Transgenic plants expressing a transgene encoding a dominant negative
15 gene product of an A-type response regulator are assayed in artificial environments or in the field to demonstrate that the transgene increases yield, delays senescence, increases photosynthesis, or increases leaf formation on the plant expressing the gene. Dominant negative transgenes are constructed according to methods known in the art. Typically, a dominant negative gene
20 encodes a mutant A-type response regulator which, when overexpressed, disrupts the activity of the wild type polypeptide.

Mutants

Plants having decreased expression of an A-type response regulator are also generated using standard mutagenesis methodologies. Such mutagenesis
25 methods include, without limitation, treatment of seeds with ethyl methylsulfate, ethylmethylsulfonate (EMS), or fast neutron irradiation, as well as T-DNA insertion methodologies. Expression of an A-type response regulator and increased yield phenotypes in mutated and non-mutated lines are evaluated according to standard procedures (for example, those described herein). Mutated

plants having decreased expression of a gene encoding a A-type response regulator exhibit increased yield relative to their non-mutated counterparts.

Use

- 5 The invention described herein is useful for a variety of agricultural and commercial purposes including, but not limited to, improving and enhancing photosynthesis, promoting cell proliferation, shoot meristem formation, promoting leaf development, increasing crop yields, improving crop and ornamental quality, and reducing agricultural production costs. In particular,
- 10 ectopic expression of a B-type response regulator (such as ARR2 or an ARR2 homolog) in a plant cell provides a method for increasing plant productivity, photosynthesis, and delaying senescence. The invention is especially useful for crop plants such as tomato, potato, cotton, pepper, maize, wheat, rice, and legumes such as soybean.
- 15 All publications and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each independent publication or patent application was specifically and individually indicated to be incorporated by reference.
- 20 What is claimed is:

CLAIMS

1. A method for increasing yield in a plant, said method comprising the steps of:
- 5 (a) introducing into plant cells a transgene comprising DNA encoding a B-type response regulator operably linked to a promoter functional in plant cells to yield transformed plant cells; and
- (b) regenerating a plant from said transformed cells, wherein said B-type response regulator is expressed in the cells of said transgenic plant, thereby increasing yield in said plant.
- 10
2. The method of claim 1, wherein said B-type response regulator is a crucifer B-type response regulator.
3. The method of claim 2, wherein said crucifer B-type response regulator
- 15 is selected from the group consisting of ARR1, ARR2, and ARR10.
4. The method of claim 1, wherein said DNA encoding said B-type response regulator is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under
- 20 cycling conditions.
5. A method for increasing yield in a plant, said method comprising the steps of:
- (a) introducing into plant cells a transgene operably linked to a promoter
- 25 functional in plant cells to yield transformed plant cells; and
- (b) regenerating a plant from said transformed cells, wherein expression of said transgene reduces expression of an A-type response regulator in the cells of said plant, thereby increasing yield in said plant.

6. The method of claim 5, wherein said A-type response regulator is a crucifer A-type response regulator.

7. The method of claim 6, wherein said crucifer A-type response
5 regulator is selected from the group consisting of ARR4, ARR5, ARR6, and ARR7.

8. The method of claim 5, wherein said transgene expresses antisense A-type response regulator RNA.

10

9. The method of claim 5, wherein said transgene expresses a dominant negative A-type response regulator.

10. The method of claim 5, wherein said transgene co-suppresses
15 expression of A-type response regulator.

11. The method of claim 5, wherein said DNA encoding said transgene is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under cycling conditions.

20

12. A method for increasing shoot formation in a plant, said method comprising the steps of:

(a) introducing into plant cells a transgene comprising DNA encoding a B-type response regulator operably linked to a promoter functional in plant cells to
25 yield transformed plant cells; and

(b) regenerating a plant from said transformed cells, wherein said B-type response regulator is expressed in the cells of said plant, thereby increasing shoot formation in said plant.

13. The method of claim 12, wherein said B-type response regulator is a crucifer B-type response regulator.

14. The method of claim 13, wherein said crucifer B-type response
5 regulator is selected from the group consisting of ARR1, ARR2, and ARR10.

15. The method of claim 12, wherein said DNA encoding said B-type
response regulator is constitutively expressed, inducibly expressed, expressed in a
cell-specific, tissue-specific, or organ-specific manner, or expressed under
10 cycling conditions.

16. A method for increasing shoot formation in a plant, said method
comprising the steps of:

(a) introducing into plant cells a transgene operably linked to a promoter
15 functional in plant cells to yield transformed plant cells; and

(b) regenerating a plant from said transformed cells, wherein expression of
said transgene reduces expression of an A-type response regulator in the cells of
said plant, thereby increasing shoot formation in said plant.

20 17. The method of claim 16, wherein said A-type response regulator is a
crucifer A-type response regulator.

18. The method of claim 17, wherein said crucifer A-type response
regulator is selected from the group consisting of ARR4, ARR5, ARR6, and
25 ARR7.

19. The method of claim 16, wherein said transgene expresses antisense
A-type response regulator RNA.

20. The method of claim 16, wherein said transgene expresses a dominant negative A-type response regulator.

21. The method of claim 16, wherein said transgene co-suppresses
5 expression of A-type response regulator.

22. The method of claim 16, wherein said transgene is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under cycling conditions.
10

23. A method for delaying senescence in a plant, said method comprising the steps of:

(a) introducing into plant cells a transgene comprising DNA encoding a B-type response regulator operably linked to a promoter functional in plant cells to
15 yield transformed plant cells; and

(b) regenerating a plant from said transformed cells, wherein said B-type response regulator is expressed in the cells of said plant, thereby delaying senescence in said plant.

20 24. The method of claim 23, wherein said B-type response regulator is a crucifer B-type response regulator.

25 25. The method of claim 24, wherein said crucifer B-type response regulator is selected from the group consisting of ARR1, ARR2, and ARR10.

26. The method of claim 23, wherein said DNA encoding said B-type response regulator is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under cycling conditions.

30

27. A method for delaying senescence in a plant, said method comprising the steps of:

(a) introducing into plant cells a transgene operably linked to a promoter functional in plant cells to yield transformed plant cells; and

5 (b) regenerating a plant from said transformed cells, wherein expression of said transgene reduces expression of an A-type response regulator in the cells of said plant, thereby delaying senescence in said plant.

28. The method of claim 27, wherein said A-type response regulator is a
10 crucifer A-type response regulator.

29. The method of claim 28, wherein said crucifer A-type response regulator is selected from the group consisting of ARR4, ARR5, ARR6, and ARR7.
15

30. The method of claim 27, wherein said transgene expresses antisense A-type response regulator RNA.

31. The method of claim 27, wherein said transgene expresses a dominant
20 negative A-type response regulator.

32. The method of claim 27, wherein said transgene co-suppresses expression of A-type response regulator.

25 33. The method of claim 27, wherein said transgene is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under cycling conditions.

34. A method for increasing yield in a plant, said method comprising the steps of:

(a) introducing into plant cells a transgene comprising DNA encoding a histidine kinase operably linked to a promoter functional in plant cells to yield transformed plant cells; and

(b) regenerating a plant from said transformed cells, wherein histidine kinase is expressed in the cells of said plant, thereby increasing yield in said plant.

35. The method of claim 34, wherein said histidine kinase is a crucifer histidine kinase.

36. The method of claim 35, wherein said crucifer histidine kinase is CKI1 or CRE1.

37. The method of claim 34, wherein said DNA encoding said histidine kinase is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under cycling conditions.

38. A method for increasing shoot formation in a plant, said method comprising the steps of:

(a) introducing into plant cells a transgene comprising DNA encoding a histidine kinase operably linked to a promoter functional in plant cells to yield transformed plant cells; and

(b) regenerating a plant from said transformed cells, wherein histidine kinase is expressed in the cells of said plant, thereby increasing shoot formation in said plant.

39. The method of claim 38, wherein said histidine kinase is a crucifer histidine kinase.

40. The method of claim 39, wherein said crucifer histidine kinase is
5 CKI1 or CRE1.

41. The method of claim 38, wherein said DNA encoding said histidine kinase is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under cycling
10 conditions.

42. A method for delaying senescence in a plant, said method comprising the steps of:

(a) introducing into plant cells a transgene comprising DNA encoding a
15 histidine kinase operably linked to a promoter functional in plant cells to yield transformed plant cells; and

(b) regenerating a plant from said transformed cells, wherein histidine kinase is expressed in the cells of said plant, thereby delaying senescence in said plant.
20

43. The method of claim 42, wherein said histidine kinase is a crucifer histidine kinase.

44. The method of claim 43, wherein said crucifer histidine kinase is
25 CKI1 or CRE1.

45. The method of claim 42, wherein said DNA encoding said histidine kinase is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under cycling
30 conditions.

46. A plant or plant component comprising at least one transgene encoding (i) an A-type response regulator polypeptide, (ii) an antisense A-type response regulator RNA, (iii) a dominant negative A-type response regulator polypeptide, (iv) a B-type response regulator polypeptide, (v) an antisense B-type response regulator RNA, (vi) a dominant negative B-type response regulator, (vii) an antisense HK RNA, (viii) a dominant negative HK polypeptide, or any combination of (i) –(viii).

47. The plant or plant component of claim 48, further comprising a transgene encoding a histidine kinase polypeptide.

48. A plant or plant component comprising:

(a) a first transgene encoding (i) an A-type response regulator polypeptide, (ii) an antisense A-type response regulator RNA, or (iii) a dominant negative A-type response regulator polypeptide or combination thereof;

(b) a second transgene encoding (iv) a B-type response regulator polypeptide, (v) an antisense B-type response regulator RNA, (vi) a dominant negative B-type response regulator or combination thereof, and

(c) a third transgene encoding (vii) a HK polypeptide; (viii) an antisense HK RNA, (ix) a dominant negative HK polypeptide or combination thereof.

49. The plant or plant component of claims 46, 47, or 48, wherein said plant or plant component is selected from the group consisting of wheat, rice, maize, barley, potato, soybean, tomato, oats, cotton, and sunflower.

FIG. 1A




FIG. 1B




FIG. 1C




FIG. 1D





FIG. 1E



4/9

FIG. 1F




FIG. 2A





FIG. 2B



6/9

FIG. 3A




FIG. 3B




FIG. 3C




FIG. 4A




FIG. 4B




FIG. 4C





FIG. 5



SEQUENCE LISTING

<110> The General Hospital Corporation

<120> Cytokinin Response Regulators and Uses
Thereof

<130> 00786/402WO2

<150> US 60/296,554

<151> 2001-06-06

<160> 30

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 2401

<212> DNA

<213> Arabidopsis thaliana

<400> 1

```

tagatcaagt tttacaactt tttttacgta cttaaaagat tgataaaatt gtacttggtg 60
ttggtgacca ttgaaacatt tccctttatt cactaaaact agcttttgca acctttttat 120
tcaactgtat tgggttggtg tatgttttta ttattcaact ctatttaatt attcatgtag 180
cttaacataa caataataac taacatgcca taaattccga agaagaaaaa ttgttataaa 240
agaactatat gcacctaaca aaaaaccaac aaaccatag tattttcttt ttgggttaca 300
taaaaatatc tttgtatgaa gaagaagaaa gaatcattac acagctttca ctcacaagat 360
tacaagtttg caaattgcat ataaggaaac tcccaaaaaa tatcatcaaa gatcttttca 420
gaacacaaaa aaaaaaagggt aaaagtatat ctcttttgcc atagttagac tcaaaaatac 480
gacatcgttt acctcacacg tgcacactca ccgacttaca gaagaacca cactacgtt 540
gagggtactt tagtcattta gcatcgaacg aaattgtctg agcggagaag aggagtttcc 600
gttatccctt tatatatatt ctctctcacc tttcttctct ttttctttga tttttattaa 660
atcaacaaaa atagaaaaaa aaacataaaa ataaaaataa aaaatcttca cgtttcttct 720
ctctctctct ctctctctct cgagccacca aatctgaatt aggggttttg agaattattca 780
tcttttgatt tcaaattctt caccactgt gtaatttcac tcgtcaggat tcatctgcgg 840
aatcatgatt acagattcga tcaccaacgc ttctgctact tcagctccga gagattccgg 900
aaagaagaag agggtaacgt tatectcttt gataaatctc attcctttct ctgaaattga 960
ttcaaagtth tgatttttaa tgggtttgtt attgcattac gttttgcaga acaataagtc 1020
ggctaagatg aagcagaaca agcttggtct ccgtcgtgag caatggcttt ctcaagggtta 1080
aaacttcate tttctctect ttaatgtttc aattctgcct gattcgtttg gtttagatttg 1140
gttgctctat tattgatttt ggggttttga ttttgatatt agttgcggtg agcaataagg 1200
aagttaaaga ggagaggagt gttaatecga gtcaaaagcc tcatcatgag agttcagata 1260
aggtgcgtag agaagaggat aacaatggtg ggaataatct tcttcatcat gagagtttta 1320
tggagtcacc ttcaaatagc tctgttggtg gtacatattc gagcactaac ttcagtggga 1380
gaagtagcag gagtagtagt agcagcagtg gcttttgctc tggtaatata acagaagagg 1440
aaaatgtaga cgatgatgat gatgggtgtg tggatgattg ggaagctgtt gctgatgcgt 1500
tagcggctga ggaagagatt gagaaaaaga gtctctctct tgagtctgtg aaagagcaag 1560
tgagtgttgg acaatcagct tetaatgtgt gtgattcgtc gattagtgat gcatcagatg 1620
ttgtgggtgt tgaagatcca aagcaggaat gcttgagagt gtcatacaag aagcagacta 1680
gtaatagagc ttggaggcta gatgatgacc ttcgcccaca ggggttacct aatttggcga 1740
agcagcttag ttttccggag ttagacaagc gttttagctc tgtggcgatt ccgtcttcat 1800
gtcccatatg ctacgaagac ttggacttga cggattcgaa tttcctcccc tgtccttggt 1860
gatttcggct ctgtctgttc tgccacaaga ccatttgcca tggagatggg cgttgtccag 1920
gctgcaggaa accctatgaa cggaatatgg tcaaggctga gactagtatt caaggtggtg 1980
gtctaacaat tcggttggct cgttcgtcta gcatgttttg caagtthtaa aaggagagg 2040
gcggttttct caaccatggt gtcttttggg actcgagaac ttaagctctg ttttctatgt 2100

```

```

catctatggt tctaagtctg aaacactgtg gtgatgatgt agaatgtgat gtgtgaatac 2160
ataaaagggtg gtacagaaaa tgattcaaata acatttagat agtttcaata atgaatgcta 2220
tgttctcttt tctaattcca tatgtttggt ctgcatttat tccttgtcaa acattattga 2280
aggtttaaga gttattttgt tgctatgggtg aatcctcttg acaagttact catgaaccaa 2340
agcttggttt ttagaatcac cattcaccag agatcaactc tcattacttc aaattctttt 2400
a 2401

```

<210> 2

<211> 1863

<212> DNA

<213> *Arabidopsis thaliana*

<400> 2

```

atggcattgt ctctgctccg gaagaacaaa catggattcg atatagtaat cagtgatgtt 60
catatgcctg acatggacgg tttcaagctt cttgagcatg ttgggtctaga gatggactta 120
cctgttatca tgatgtctgc ggatgattca aagagtgtgg ttctaaaggg agtaacgcac 180
ggtgcggttg attaccttat caagcctgta cgtatggagg cacttaagaa catatggcag 240
catgtagtta ggaagaggag aagtgaatgg agtgtaccgg aacattctgg gagcattgag 300
gagactggcg agagacagca gcagcaacat agaggagggtg gtggtggtgc agctgtttct 360
ggtggagagg atgcggtgga tgataactca tcctcggtta acgaaggtaa caattggagg 420
agcagttcac ggaagaggaa agacgaggaa ggagaagagc aaggagatga taaggacgaa 480
gatgcgtcga atttgaagaa accgcgtgtc gtctggtctg ttgaattgca tcagcagttt 540
gttgctgctg ttaatcagct cggcgttgag aaggcgggtc ctaaaaagat cttagagctg 600
atgaatgttc ctggtctaac ccgagaaaac gtagcaagtc acctccagaa ataccggata 660
tatctaagac ggcttgaggg ggtatcgtag caccaaggca atcttaacaa ctcgtttatg 720
acgggtcagg atgcgagctt cggacctctt tcgacattga atggggttga tcttcaagca 780
ctagccgtca caggtcagtt acctgcacag agtcttgcac agcttcaagc cgctggttta 840
ggccggcctg cgatggtctc taagtcaggt ttgcccgttt cctccattgt ggatgagaga 900
agcatcttca gctttgacaa cacgaaaaca agatttgtag aagggttgg gcatcacggg 960
caacaacccc aacagcaacc acagatgaac ttacttcacg gtgtccccac ggggtttacaa 1020
cagcagcttc ctatgggtaa tcgaatgagt attcaacaac agattgctgc tgttcgagct 1080
ggaaatagtg ttcaaaacaa cggaatgctg atgcctctag cgggtcagca gtctttgcct 1140
cggggaccac cgcctatgct aacctcttcg caatcatcca tcaggcagcc gatgttatca 1200
aaccgcattt ccgagagaag tggtttctct ggaaggaaca atatccccga gagcagcaga 1260
gtgttaccca caagttacac taatctcaca acacaacact catcaagctc gatgccttat 1320
aacaacttcc aaccagaact tcccgtgaac agtttcccgc tggcaagtgc accagggata 1380
tcagtaccgg ttccgaaagc cacttcttac caggaagagg ttaacagctc cgaagcgggt 1440
ttcactacgc cgagctacga catgttcacc accagacaga atgattggga tctgaggaat 1500
attggaatag cctttgactc acatcaggac tcagaatccg ctgcgttttc cgcttcagaa 1560
gcctactctt cttcgtccat gtcaagacac aacacgacag ttgcagccac cgagcatggc 1620
cgaaaccacc agcagccacc atcgggaatg gtacagcacc atcaggttta tgcagacgga 1680
aacgggtggt cagtgagggt gaaatcagag agagtggcta cggatacagc aacaatggcg 1740
tttcacgagc agtatagtaa tcaagaagat cttatgagcg cacttcttaa gcaggaaggg 1800
attgcaccgg ttgatggcga attcgacttt gacgcatact ccatcgataa cattccggtt 1860
tga 1863

```

<210> 3

<211> 1935

<212> DNA

<213> *Arabidopsis thaliana*

<400> 3

```

atggtaaatc cgggtcacgg aagaggaccc gattcgggta ctgctgctgg tgggtcaaac 60
tccgaccctg ttcttgcgaa tcttcgagtt cttgtcgttg atgatgatcc aacttgtctc 120
atgatcttag agaggatgct tatgacttgt ctctacagag agcagagagc gcattgtctc 180
tgcttcggaa gaacaaagaa tggttttgat attgtcatta gtgatgttca tatgcctgac 240
atggatggtt tcaagctcct tgaacacggt ggtttagaga tggatttacc tggtatcaat 300
ctgaatgttt tgaaaccttt gggttatagt atgtctgcgg atgattcgaa gagcgttgtg 360

```

```

ttgaaaggag tgactcacgg tgcagttgat tacctcatca aaccgggtacg tattgaggct 420
ttgaagaata tatggcaaca tgtggtgcgg aagaagcgta acgagtggaa tgtttctgaa 480
cattctggag gaagtattga agatactggc ggtgacaggg acaggcagca gcagcatagg 540
gaggatgctg ataacaactc gtcttcagtt aatgaaggga acgggaggag ctcgaggaag 600
cggaagggaag aggaagtaga tgatcaaggg gatgataagg aagactcatc gagtttaaag 660
aaaccacgcg tggtttggtc tgttgaattg catcagcagt ttgttgctgc tgtgaatcag 720
ctaggcggtg acagtgaagt aaaaacttgc ttgcttatgc atttgtgtgt gtcgattggt 780
aacattgtgg aattccagaa gtatcggata tatctgagac ggcttggagg agtatcgcaa 840
caccaaggaa atatgaacca ttcgtttatg actgggtcaag atcagagttt tggacctctt 900
tcttcggttg atggatttga tcttcaatct ttagctgtta ctggtcagct ccctcctcag 960
agccttgcac agcttcaagc agctggtctt ggccggccta cactcgctaa accagggatg 1020
tcggtttctc cccttgtaga tcagagaagc atcttcaact ttgaaaaccc aaaaataaga 1080
tttgagagac gacatggtca gacgatgaac aatggaaatt tgcttcattg tgtcccaacg 1140
ggtagtcaca tgcgtctgcg tcctggacag aatgttcaga gcagcggaat gatgttgcca 1200
gtagcagacc agctacctcg aggaggacca tcgatgctac catccctcgg gcaacagccg 1260
atattgtcaa gcagcgtttc aagaagaagc gatctcactg gtgcgctggc ggtagaaac 1320
agtatccccg agaccaacag cagagtgtta ccaactactc actcggctct caataacttc 1380
cccgcggatc tacctcgagc cagcttcccg ttggcaagtg ccccgaggat ttcagttcca 1440
gtatcagttt cttaccaaga agaggtcaac agctcggatg caaaaggagg ttcattcagct 1500
gctactgctg gatttggtaa cccaagctac gacatattta acgattttcc gcagcaccaa 1560
cagcacaaca agaacatcag caataaacta aacgattggg atctgcggaa tatgggattg 1620
gtcttcagtt ccaatcagga cgcagcaact gcaaccgcaa ccgcagcatt ttccacttcg 1680
gaagcatact cttcgtcttc tacgcagaga aaaagacggg aaacggacgc aacagttgtg 1740
ggtgagcatg ggcagaacct gcagtcaccg agccggaatc tgtatcatct gaaccacgtt 1800
tttatggacg gtggttcagt cagagtgaag tcagaaagag tggcggagac agtgacttgt 1860
cctccagcaa atacattgtt tcacgagcag tataatcaag aagatctgat gagcgcattt 1920
ctcaaacagg tttga 1935

```

<210> 4

<211> 599

<212> DNA

<213> Artificial Sequence

<220>

<223> derived from *Arabidopsis thaliana*

<400> 4

```

gctttgaaga atatatggca acatgtggtg cggaagaagc gtaacgagtg gaatgtttct 60
gaacattctg gaggaagtat tgaagatact ggcggtgaca gggacaggca gcagcagcat 120
agggaggatg ctgataacaa ctctgtcttc gttaatgaag ggaacgggag gagctcgagg 180
aagcgggaag aagaggaagt agatgatcaa ggggatgata aggaagactc atcagattta 240
aagaaaccac gcgtggtttg gtctgttgaa ttgcatcagc agtttgttgc tgcgtgaat 300
cagctaggcg ttgacagtga gttaaaaact tgcttgctta tgcatttgtg tgtgtcgatg 360
tttgttttct tgtctaactt ttaccgtatt ggggcattca atgcagaagc tgttcctaag 420
aagatcttag agatgatgaa tgtacccggg ctaacgcgag aaaacgtagc cagtcacctc 480
caggtatata tgtgatcgaa tagtattaca ttttttcatt gttgatgtgt tctaacagtg 540
gtaacattgt ggaattccag aagtatcgga tataatctgag acggcttgga ggagtatcg 599

```

<210> 5

<211> 780

<212> DNA

<213> *Arabidopsis thaliana*

<400> 5

```

atggccagag acggtggtgt ttcttgttta cgaaggtcgg agatgatgag cgtcgggtggt 60
atcggaggaa ttgaatctgc gccgttggat ttagatgaag ttcattgtct agccgttgat 120
gacagtctcg ttgatcgatg tgtcatcgag agattgcttc gtattacttc ctgcaaagtt 180
acggcggtag atagtggatg gcgtgctctg gaatttctag ggttagataa tgagaaagct 240

```



```

tctgctgaat tcgatagatt gaaagttgat ttgatcatca ctgattactg tatgcctgga 300
atgactgggt atgagcttct caagaagatt aaggaatcgt ccaatttcag agaagttccg 360
gttgtaatca tgcgtcggga gaatgtattg accagaatcg acagatgcct tgaggaaggt 420
gctcaagatt tcttattgaa accggtgaaa ctgcgccgac tgaaacgtct gagaagtcac 480
ttaactaaag acgttaaact ttccaacgga aacaaacgga agcttccgga agattctagt 540
tccgttaaact cttcgcttcc tccaccgtca cctccgttga ctatctcgcc tgaatcgctg 600
ccgccgttga ctgtttcgac tgaatcgctg gattcgctct cgccgttatc tccggtggag 660
atcttttcca cctcgccact ttcattctcg atagacgatg aagatgatga cgtgttgacg 720
tcgtcgtcgg aggaatcgcc gattcgacgg cagaagatga ggagtcccgg attagattag 780

```

<210> 6

<211> 891

<212> DNA

<213> *Arabidopsis thaliana*

<400> 6

```

gttgattctc tctatctctc tcacgagtcg cgatcctact cttcttgata tggctgaggt 60
tttgcgtccc gagatgttag atatctctaa cgacacttct tcattagcat caccgaaact 120
tcttcattgt cttgctgttg atgatagtat gggtgatcgg aagttcatcg agcggttact 180
cagagtctca tcttgtaaag ttactgttgt cgatagtgcg acaagagctt tacaatatct 240
tggattagat ggagagaata atagttcggg tggatttgag gatctgaaga ttaatttgat 300
aatgacggat tactctatgc ctgggatgac tggatatgaa ctactgaaga agatcaaaga 360
atcatcagct ttcagagaaa taccagttgt gattatgtct tcagagaaca tcttgccctc 420
tatcgataga tgtcttgaag aaggagctga agatttctta ttgaagcctg tgaaattagc 480
tgatgtgaag agattaagag attctttaat gaaagctgag gaaagagctt ttaaggatat 540
tatgcacaag agagagcttg aagctaata tatctactcg cagctaaaac gcgcaaagat 600
ctgagtttat cagattccac gtacgatatg ttcttttaaa gctcaaagat tcacacacat 660
gcatgtcctg attctttcgg cttacaattt ttgagacatt acggttctt ttgctgatag 720
aaccaagact gatcaagagt gacaaagatg gggtttttata agatagaaat gtaaattgta 780
gacttggatt cattagagag gagaagagat acattttttt tttctttaca tgagagatat 840
gtaaaatggg ggtgtgttat tttggaagtt aacatatgaa tttcttttgt t 891

```

<210> 7

<211> 942

<212> DNA

<213> *Arabidopsis thaliana*

<400> 7

```

atcctccaac attcgttgat caatggctga agttatgcta ccgaggaaga tggagattct 60
caaccattct tcaaagtttg gatcaccgga tcctcttcat gttcttgccg tcgacgacag 120
tcacgttgat cgtaaattca tcgagcgttt gctcagagta tcttcttgca aagttactgt 180
tggtgatagt gcaacaagag ctctccaata ccttggtttg gatgttgagg aaaaatcagt 240
cggttttgag gatttgaagg ttaatttaat tatgactgat tactcaatgc ccggtatgac 300
tggatatgaa ctcttgaaga agatcaaaga atcctcagct tttagagaag taccggtggg 360
aattatgtcc tccgagaaca ttttgccctg tattgataga tgtcttgaag aaggggctga 420
agatttctta ctgaagccgg taaaactctc ggatgtaaaa agattaagag attctctaata 480
gaagggttgaa gatttgtctt tcacaaagag tattcagaag agagagctag aaacagagaa 540
tgtctaccct gttcactcgc agctcaaacg cgcaaagatc tgagctctcc gatgcaaatt 600
ccgtgactgg atcttagaaa aagccttttc atatacagga agcttcaaga ctcaatgcac 660
ttgagctaca ctgatgatc tcgccagctt aatgacctat acaattttcc aaccctatg 720
gcctttttcg ctgattgaat tcagtgggtt tattcatata gtagtaatgt aaatggtata 780
gagatgagca aaagctatat tatatattat attcattaga ggggtcatag cttttttcct 840
tacaagagat ttgtaagctg gcttaataga agttgttaag aagcctcact ttatgtgaat 900
aaatgatttt gttagctaata cataaacaat cagaagtttc ct 942

```

<210> 8

<211> 740

<212> DNA

<213> *Arabidopsis thaliana*

<400> 8

```

caatctgctc tcttttttat tctgagtttg acaatggcgg ttggtgaggt catgaggatg 60
gagattcccg ccggtggaga tttgactgtt actactccgg agttgcatgt tcttgccgtc 120
gatgatagta ttgtggatcg taaagtcac gagaggttgc ttagaatctc ttcttgtaaa 180
gtgacgactg tagagagtgg aactagggct ttgcagtatc ttggcttaga tggaggcaaa 240
ggggcttcta atcttaagga tttgaagggtg aatttgatag tgacggatta ctcaatgcca 300
ggactttcag gatatgatct ccttaaaaag attaaggaat cttcagcatt cagagaagta 360
ccagtagtga ttatgtcac tgagaacatc ttacctcgta tacaagaatg tctcaaagaa 420
ggagcagagg aattcttggt gaaaccgggtg aagctagcag atgtaaagcg aatcaaaca 480
cttataatga ggaatgaagc tgaggaatgc aaaatcttaa gccattctaa caagagaaag 540
cttcaagaag acagtgatac atcatcatca agtcatgatg atacttctat caaggactct 600
tcatgttcaa aacgaatgaa atcagaatct gaaaaccttt tttctctact ttgaattata 660
tgtatagagg ccaagacctt agcaaaacca ttctgttaat attcctcaag aaagacaatg 720
tttgagagag tcatcacact

```

<210> 9

<211> 1659

<212> DNA

<213> *Arabidopsis thaliana*

<400> 9

```

atgactatgg agcaagaaat tgaagtcttg gaccagtttc cggttgggat gagagttctt 60
gctgttgacg atgaccagac ttgtctccgt attctccaga ctttgcttca gcgctgccaa 120
tatcacgtta caacaacgaa tcaggcacag accgcattgg agttgttgag ggagaacaag 180
aataagtttg atcttggtat tagcgatgtc gacatgccag acatggatgg tttcaagctg 240
cttgagcttg ttggtcttga aatggactta cctgtcataa tggtatctgc gcatagcgat 300
ccaaagtatg tgatgaaagg agtcaagcac ggtgcctgtg attatctgct taaaccgggt 360
cgaattgagg agcttaagaa catatggcaa catgttgtga gaaagagcaa acttaagaag 420
aataagagca atgtgagtaa tggttcagga aactgtgata aagcaaacag aaaacgtaaa 480
gaacagtatg aagaggagga agaggaagaa agagggaaatg ataagatga tccaacggcg 540
cagaagaagc ctctgttctt ttggacgcac gagctgcaca ataaattcct agcagctgtt 600
gatcatttag gcgttgagag agctgttcca aaaaagattc tagatctgat gaatgttgac 660
aaactcacta gagagaatgt tgcaagccac cttcagaaat tccgcgttgc tctgaagaag 720
gtgtctgatg acgccattca acaagctaac agggcggcta ttgactcaca ttttatgcaa 780
atgaattctc agaaaggact tgggtgcttc taccaccacc accgcggaat acctgttgga 840
tccggtcagt tccatgggtg aaccacaatg atgaggcatt actcttcaaa taggaatctt 900
ggctcgtctga attcccttgg agcaggaatg ttccaaccag tctcatcatc gtttctctgt 960
aaccataatg atggaggaaa catacttcag ggtttgccgc tagaagagct tcagatcaac 1020
aacaacatca acagggcttt tccaagcttt acttcacaac aaaactctcc aatggtagct 1080
cccagtaatc tgttacttct cgagggtaac ccgcagtcac catctttacc ctcaaaccgg 1140
ggtttttctc ctcatcttga gatcagcaag cgtctagaac attggtcaaa cgctgcattg 1200
tcaaccaaca ttccacagag tgatgttcat tcaaaacctg acaccttga atggaatgcg 1260
ttctgcgact cagctagtcc gctagtaaac ccaaacctgg atacaaatcc ggcattcttc 1320
tgcagaaaca cgggttttgg atccacaaat gctgcacaaa cagacttctt ttatccatta 1380
cagatgaatc agcagcctgc aaacaactca ggtccagtga cagaagctca actgtttaga 1440
agtagcaatc caaacgaagg tttactcatg ggacaacaga agcttcagag tggtttgatg 1500
gcttctgatg ctggttcctt agatgatata gtcaattcct taatgacaca ggaacagagc 1560
caatctgatt tctcggaagg tgattgggat ttggatgggt tagctcactc ggaacatgca 1620
tacgagaaac tccattttcc cttttctttg tcagcttga

```

<210> 10

<211> 1935

<212> DNA

<213> *Arabidopsis thaliana*

<400> 10

```

atggtaaatac cgggtcacgg aagaggaccc gattcgggta ctgctgctgg tgggtcaaac 60
tccgaccctgt ttcctgcgaa tcttcgagtt cttgtcgttg atgatgatcc aacttgtctc 120
atgatcttag agaggatgct tatgacttgt ctctacagag agcagagagc gcatttgtctc 180
tgcttcggaa gaacaaagaa tggttttgat attgtcatta gtaatgttca tatgcctgac 240
atggatgggtt tcaagctcct tgaacacggt ggttttagaga tggatttacc tgttatcaat 300
ctgaatgttt tgaaaccttt ggttatagtg atgtctgcgg atgattcgaa gagcgttgtg 360
ttgaaaggag tgactcacgg tgcagttgat tacctcatca aacgggtacg tattgaggct 420
ttgaagaata tatggcaaca tgtggtgcgg aagaagcgta acgagtggaa tgtttctgaa 480
cattctggag gaagtattga agatactggc ggtgacaggg acaggcagca gcagcatagg 540
gaggatgctg ataacaactc gtcttcagtt aatgaaggga acgggaggag ctcgaggag 600
cggaaggag aggaagtaga tgatcaaggg gatgataagg aagactcatc gagtttaaag 660
aaaccacgcg tggtttggtc tgttgaattg catcagcagt ttgttgctgc tgtgaatcag 720
ctaggcgttg acagtgagtt aaaaacttgc ttgcttatgc atttgtgtgt gtcgattggg 780
aacatttgtg aattccagaa gtatcggata tatctgagac ggcttgagg agtatcgcaa 840
caccaaggaa atatgaacca ttcgtttatg actggtcaag atcagagttt tggacctctt 900
tcttcgttga atggatttga tcttcaatct ttagctgtta ctggtcagct cctcctcag 960
agccttgac agcttcaagc agctggtctt ggccggccta cactcgctaa accagggatg 1020
tcggtttctc ccttgtaga tcagagaagc atcttcaact ttgaaaaccc aaaaataaga 1080
tttgagacg gacatggtca gacgatgaac aatggaaatt tgcttcatgg tgtcccaacg 1140
ggtagtcaca tgcgtctgcg tcctggacag aatgttcaga gcagcggaat gatgttgcca 1200
gtagcagacc agctacctcg aggaggacca tcgatgctac catcctcgg gcaacagccg 1260
atattgtcaa gcagcgttcc aagaagaagc gatctcactg gtgcgctggc ggtagaaac 1320
agtatcccc agaccaacag cagagtgtta ccaactactc actcggctct caataacttc 1380
cccgcggatc tacctcgcag cagcttccc ttggcaagtg ccccgaggat ttcagttcca 1440
gtatcagttt cttaccaaga agaggtcaac agctcggatg caaaaggagg ttcacagct 1500
gctactgctg gatttggtta ccaagctac gacatattha acgattttcc gcagcaccaa 1560
cagcacaaca agaacatcag caataaacta aacgattggg atctgcggaa tatgggattg 1620
gtcttcagtt ccaatcagga cgcagcaact gcaaccgcaa ccgcagcatt ttccacttcg 1680
gaagcatact cttcgtcttc tacgcagaga aaaagacggg aaacggacgc aacagtttgt 1740
ggtgagcatg ggcagaacct gcagtcaccg agccggaatc tgtatcatct gaaccacgtt 1800
tttatggacg gtggttcagt cagagtgaag tcagaaagag tggcgagagc agtgacttgt 1860
cctccagcaa atacattgtt tcacgagcag tataatcaag aagatctgat gagcgcattt 1920
ctcaaacagg tttga

```

<210> 11

<211> 780

<212> DNA

<213> Artificial Sequence

<220>

<223> derived from *Arabidopsis thaliana*

<400> 11

```

atggccagag acggtggtgt ttcttgttta cgaaggtcgg agatgatgag cgtcgggtgg 60
atcgaggagaa ttgaatctgc gccgttggtt ttagatgaag ttcattgtct agccgttgat 120
gacagtctcg ttgatcgtat tgtcatcgag agattgcttc gtattacttc ctgcaaagtt 180
acggcggtag atagtggatg gcgtgctctg gaatttctag ggtagataaa tgagaaagct 240
tctgctgaat tcgatagatt gaaagttgat ttgatcatca ctaattactg tatgcctgga 300
atgactgggt atgagcttct caagaagatt aaggaatcgt ccaatttcag agaagttccg 360
gttgtaatca tgtcgtcgga gaatgtattg accagaatcg acagatgcct tgaggaagg 420
gctcaagatt tcttattgaa accggtgaaa ctcccgacg tgaaacgtct gagaagtcac 480
ttaactaaag acgttaaact ttccaacgga aacaaacgga agcttccgga agattctagt 540
tccgttaact cttcgtcttc tccaccgtca cctccgttga ctatctcgcc tgaatcgtcg 600
ccgccgttga ctgtttcgac tgaatcgtcg gattcgtctc cgccgttatc tccggtggag 660
atcttttcca cctcgccact ttcattctcg atagacgatg aagatgatga cgtgttgacg 720
tcgtcgtcgg aggaatcgcc gattcgacgg cagaagatga ggagtcgagg attagattag 780

```

<210> 12
 <211> 942
 <212> DNA
 <213> Arabidopsis thaliana

<400> 12
 atcctccaac attcgttgat caatggctga agttatgcta ccgaggaaga tggagattct 60
 caaccattct tcaaagtttg gatcaccgga tctcttcat gttcttgccg tcgacgacag 120
 tcacgttgat cgtaaattca tcgagcggtt gtcagagta tcttcttgca aagttactgt 180
 tgttgatagt gcaacaagag ctctccaata ccttggtttg gatgttgagg aaaaatcagt 240
 cggttttgag gatttgaagg ttaatttaat tatgactaat tactcaatgc ccggtatgac 300
 tggatatgaa ctcttgaaga agatcaaaga atcctcagct tttagagaag taccggtggt 360
 aattatgtcc tccgagaaca ttttgcctcg tattgataga tgtcttgaag aaggggctga 420
 agatttctta ctgaagccgg taaaactctc ggatgtaaaa agattaagag attctctaata 480
 gaagggttgaa gatttgtctt tcacaaagag tattcagaag agagagctag aaacagagaa 540
 tgtctacctt gttcactcgc agctcaaagc cgcaaagatc tgagctctcc gatgcaaatt 600
 ccgtgactgg atcttagaaa aagccttttc atatacagga agcttcaaga ctcaatgcac 660
 ttgagctaca ctgatgattc tcgccagctt aatgacctat acaattttcc aaccttatg 720
 gccttttttcg ctgattgaat tcagtgggtt tattcatata gtagtaatgt aaatgggtata 780
 gagatgagca aaagctatat tatatattat attcattaga ggggtcatag cttttttcct 840
 tacaagagat ttgtaagctg gcttaataga agttgttaag aagcctcact ttatgtgaat 900
 aatgatttt gttagctaat cataaacaat cagaagtttc ct 942

<210> 13
 <211> 3677
 <212> DNA
 <213> Arabidopsis thaliana

<400> 13
 aattattcac tcaaattcac aaagggtttc gactttgctc cgaggaagaa gataatatga 60
 aaagagcttt ttagggttta tcattctcct tgactttgca aaacgtgaaa taagcttctt 120
 ccttgaggga ctaatcaaga cagaaatctg ttcctctaaa aacgatcgcc gttctagtct 180
 actcaaata tgggtgaaagt tacaagctt gtggcttcac gtccaattgt ggtcttttgc 240
 gtcttggeat tcttggtggt tgttttcgag tgcatttgga tctcaaattg gcgaacaaca 300
 acggagaacc tagtcaaaga ggctgcttca tttaccgaag atctccggac aagtctagtt 360
 tcggagattg aaaacatcgg aaaatttaca tatgctaaga caaacttata tacgatcgg 420
 ttagcgagag ttatagattc ttatatcacc aacaacgaca ctggttttac agagattcaa 480
 acacagatcg caccattggt gtttgtagct tattcaacga tccttcaagt ctcaacagtt 540
 tcgtacatca gtagggacgg tctcatgttt tcttacattg cagaatcaaa cacaagtgtc 600
 gctgtttttg ccaattcttc gtcgaattca agtcgtggag actacacttg gtacactcaa 660
 accgtggatc agttaactgg tcgtcttaac gggaaactca cgaaatctca gtcgttagat 720
 gtaaccata cagattgggt ccaagcagca cagagtaata actacactac agcctttgta 780
 ggaacgagct tgggaggaga agataacgag actctaatac agagcgtgggt tagcttgtac 840
 agcaagaaag gtcttggttc tttagggttt ccggttaaga ctttaaccga agttttgaac 900
 agtttgaatc tacacggcga agagctttac atgtggacaa aggacgggac ggtgcttgtt 960
 cgtgaagggt cactgaatga tcttttcttc atctccaatg gctcgatttg cttcggtaga 1020
 gaatcgaact ccctctgggt tcaatgcate cctgaaaatt gcagttccag tggctacgag 1080
 gtggagatca aaagattaa gataccaagct ttttgctctg ttattgaagt ttcgggctgt 1140
 cctctgagat acacactcat gtttcccaac aaaggaggag caacacgcat caagcaccaa 1200
 gcggaaaagg caaaatatca acttattgtg gttatgatat ttcttggtct cggttggcct 1260
 gtatggtttg tgtggtttat gatgcaagca acaaggagag agatgcata gcgtgcaacg 1320
 ctgataaacc aaatggaagc gacacaacaa gctgagagaa agagcatgaa caagagtcaa 1380
 gcatttgcaa atgctagcca cgatattaga ggtgcccttg cagggatgaa aggtctgatt 1440
 gatatatgtc gtgatggagt taaacctggc tccgacgtag acaccactct caaccaagt 1500
 aatgtttgcg ccaaggattt ggttgctctg ctcaactctg ttttggacat gagcaaaatc 1560
 gaaagcggga agatgcagtt agtgaagaa gatttcaact tgtcgaaact tcttgaagac 1620
 gtcacgatt tttaccatcc cgttgcatg aagaaagggg ttgatgtagt tttggatccg 1680
 cacgatggct cggttttcaa attctcgaat gtacgagggg atagtggcag actgaagcag 1740


```

attctgaaca atcttgtag caatgctgtc aagttcacccg tcgacgggca cattgcggta 1800
agagcttggg ctcagaggcc aggttccaat agctctgtgg tccttgcac atatectaaa 1860
ggtgtgtcca agtttgtaaa gagtatgttc tgcaagaata aagaagagtc atcaacctac 1920
gagacagaaa tatcgaatc cataagaaac aatgcaaaca cgatggagtt tgtgtttgaa 1980
gtggatgata ctggtaaagg gatacctatg gagatgcgta agtcgggtatt tgaaaactat 2040
gttcaagtaa gagaaacagc tcaaggacac caaggaactg gtttagggct cgggattgtg 2100
cagtctttgg taagattaat gggaggggag ataagaatca ccgacaaggc catgggagag 2160
aaaggaacat gtttccaatt caatgtttta ttgacaacat tagagtctcc tccagttagt 2220
gacatgaaag tgagacagga gatcgaagca ggagggcatt atgtatccac gccaaacctc 2280
gggctgacta taaacacttc acttgagggt agcatgaata tacgtaacct gagtcctaga 2340
ttcaacaact gtctcagctc aagtccaaag caagaagggt ctagagtggg tcttctattg 2400
aaaaatgaag aacgcagaag agttacagag aaatacataa agaactctgg gattaaagtt 2460
acagtgggtg agaaatggga gcatttgagt tatgctttgg agagactttt tggattttca 2520
cctcagagtt ccatgggaag agcagagtggt agtttgtcat gtccgagctc aaggaggtta 2580
cctttcattg gcatggacgg tattgattca agaagccaac ttcctaaaag gagaagcatc 2640
agtttctctg cagttgtcct tttggtgatt gatgcaaaaa ctggaccatt ttttgagctg 2700
tgcatattg tcaaacagtt tcgtagaggc ttgccccatg gaatatcctg taaagttgtt 2760
tggcttaacg aatcgagcac tcgtgtaagt gagagagggg acattagttg ttcgagacct 2820
ttgcacggat cgcgtcttat ggaagtcttg aagatgttgc ctgaatttgg aggaactgtg 2880
ctaaaagaac cacctactga gctgcaaagg gaatcattgc tgagacattc ttttgttgca 2940
gagagatcac caaaacataa agtccaagaa gagggggcaa gctcaatgtt taacaaaaaa 3000
ttaggtaaga ggataatggc atcaacagat tcagagagtg agactagggt caagtcagtg 3060
cgtaccggtc gaaagcctat tgggaaccca gaggacgagc aagagacttc caagccgagt 3120
gacgatgaat tcttaagagg aaagagagtt cttgtggtcg atgataactt tatatcacgt 3180
aaagttgcaa caggaaagct gaagaagatg ggagtctcag aggtcgaaca atgcgacagt 3240
gggaaagaag ctttgagatt agtcactgaa gggcttacac aaagagaaga acaaggttca 3300
gtagataaac ttccgtttga ctacatattc atggactgcc aaatgccaga aatggatggc 3360
tatgaagcaa ctagagagat taggaaagtg gagaaaagtt atgggggtgcg tacaccaatt 3420
atagctgtat ctggtcatga tcctgggttca gaggaagcaa gagaaacctt tcaagctgga 3480
atggacgcct tcttagataa aagcttgaat caacttgcaa acgtcattag agaaatcgaa 3540
agcaaacgtc actagtattg tataagtaga cagcacgcag ctttgtgcta ctaccattag 3600
tgttttagac aatacttttt tagtatgacg atgtcatagc agttttcttc taagtttatt 3660
taaagtccat ttcagat 3677

```

<210> 14

<211> 3677

<212> DNA

<213> Artificial Sequence

<220>

<223> derived from *Arabidopsis thaliana*

<400> 14

```

aattattcac tcaaattcac aaagggtttc gactttgtctc cgaggaagaa gataatatga 60
aaagagcttt ttagggttta teattctcct tgactttgca aaacgtgaaa taagcttctt 120
ccttggagga ctaatcaaga cagaaatctg ttctctaaa aacgatcgcc gttctagttc 180
actcaaatga tggtgaaagt taaaaagctt gtggcttcac gtccaattgt ggtcttttgc 240
gtcctggcat tcctgggtggg tgttttcgag tgcatttggg tctcaaattg gcgaacaaca 300
acggagaacc tagtcaaaga ggtcgcttca tttaccgaag atctccggac aagtctagtt 360
tcggagattg aaaacatcgg aaaatttaca tatgctaaga caaacttate tacgatcggg 420
ttagcgagag ttatagattc ttatatcacc aacaacgaca ctggttttac agagattcaa 480
acacagatcg caccattgtt gttttagact tattcaacga tccttcaagt ctcaagtt 540
tcgtacatca gtagggacgg tctcatgttt tcttacattg cagaatcaaa cacaagtgtc 600
gctgtttttg ccaattcctc gtogaattca agtcgtggag actacacttg gtacactcaa 660
accgtggatc agttaactgg tcgtcttaac gggaactcaa cgaaatctca gtcgttagat 720
gtaaccata cagattgggt ccaagcagca cagagtaata actacactac agcctttgta 780
ggaacgagct tgggaggaga agataacgag actctaatac agagcgtggg tagcttgta 840
agcaagaaag gtcttgtttc tttagggttt ccggttaaga ctttaaccga agttttgaac 900

```

```

agtttgaatc tacacggcga agagctttac atgtggacaa aggacgggac ggtgcttggt 960
cgtgaagggt cactgaatga ttctttcttc atctccaatg gctcgatttg cttcggtaga 1020
gaatcgaact ccctctggtc tcaatgcac cctgaaaatt gcagttccag tggctacgag 1080
gtggagatca aaagattaag ataccaagct ttttgcctg ttattgaagt ttcgggcgtt 1140
cctctgagat acacactcat gtttcccaac aaaggaggag caacacgcac caagcaccaa 1200
gcggaaaaagg caaaatatca acttattgtg gttatgatat ttcttggctt cggttggcct 1260
gtatggtttg tgtggtttat gatgcaagca acaaggagag agatgcatat gcgtgcaacg 1320
ctgataaacc aaatggaagc gacacaacaa gctgagagaa agagcatgaa caagagtcaa 1380
gcatttgcaa atgctagcca agatattaga ggtgcccttg cagggatgaa aggtctgatt 1440
gatatatgtc gtgatggagt taaacctggc tccgacgtag acaccactct caaccaagtg 1500
aatgtttgcg ccaaggattt ggttgcctg ctcaactctg ttttggacac gagcaaaatc 1560
gaaagcggga agatgcagtt agtggaaaga gatttcaact tgtcgaaact tcttgaagac 1620
gtcatcgatt ttaccatcc cgttgcgatg aagaaagggg ttgatgtagt tttgatccg 1680
cacgatggct cggttttcaa attctcgaat gtacgagggg atagtggcag actgaagcag 1740
attctgaaca atcttgtag caatgctgtc aagttcaccc tgcacgggca cattgcggtg 1800
agagcttggg ctcagaggcc aggttccaat agctctgtgg tccttgcac atatectaaa 1860
ggtgtgtcca agtttgtaaa gagtatgttc tgcaagaata aagaagagtc atcaacctac 1920
gagacagaaa tatcgaattc cataagaac aatgcaaaca cgatggagtt tgtgtttgaa 1980
gtggatgata ctggtaaagg gatacctatg gagatgcgta agtcggtatt tgaaaactat 2040
gttcaagtaa gagaaacagc tcaaggacac caaggaactg gtttagggct cgggattgtg 2100
cagtctttgg taagattaat gggaggggag ataagaatca ccgacaaggc catgggagag 2160
aaaggaacat gtttccaatt caatgtttta ttgacaacat tagagtctcc tccagtgagt 2220
gacatgaaag tgagacagga gatcgaagca ggaggcgatt atgtatccac gccaaacctc 2280
gggctgacta taaacacttc acttggaggt agcatgaata tacgtaacct gagtctaga 2340
ttcaacaact gtctcagctc aagtccaaag caagaaggtt cttagagtgt tcttctattg 2400
aaaaatgaag aacgcagaag agttacagag aaatacataa agaactcttg gattaaagtt 2460
acagtgggtg agaaatggga gcatttgagt tatgcttttg agagactttt tggattttca 2520
cctcagagtt ccatgggaag agcagagtggt agtttgtcat gtccgagctc aaggaggtta 2580
cctttcattg gcattggacgg tattgattca agaagccaac ttcctaaaag gagaagcatc 2640
agtttctctg cagttgtcct tttggtgatt gatgcaaaaa ctggaccatt ttttgagctg 2700
tgcgatattg tcaaacagtt tcgtagagge ttgccccatg gaatactctg taaagttgtt 2760
tggtctaacg aatcgagcac tcgtgtaagt gagagagggg acattagttg ttcgagacct 2820
ttgcacggat cgcgtcttat ggaagtcttg aagatgttgc ctgaatttgg aggaactgtg 2880
ctaaaagaac cacctactga gctgcaaagg gaatcattgc tgagacattc ttttgttgca 2940
gagagatcac caaaacataa agtccaagaa gaggggcca gctcaatgtt taacaaaaaa 3000
ttaggtaaga ggataatggc atcaacagat tcagagagtg agactagggt caagtcagtg 3060
cgtaccggtc gaaagcctat tgggaaccca gagacgagc aagagacttc caagccgagt 3120
gacgatgaat tcttaagagg aaagagagtt cttgtggtcg atgataactt tatatcacgt 3180
aaagttgcaa caggaaagct gaagaagatg ggagtctcag aggtcgaaca atgacgacgt 3240
gggaaagaag ctttgagatt agtcaactgaa gggcttacac aaagagaaga acaaggttca 3300
gtagataaac ttccgtttga ctacatatc atggactgcc aaatgccaga aatggatggc 3360
tatgaagcaa ctagagagat taggaaagtg gagaaaagtt atggggtgcg tacaccaatt 3420
atagctgtat ctggtcatga tcctggttca gaggaagcaa gagaaacat tcaagctgga 3480
atggacgcct tcttagataa aagcttgaat caacttgcaa acgtcattag agaaatcgaa 3540
agcaaacgtc actagtattg tataagtaga cagcacgcag ctttgtgcta ctaccattag 3600
tgtttttagac aatacttttt tagtatgacg atgtcatagc agttttcttc taagtttatt 3660
taaagtccat ttcagat 3677

```

<210> 15

<211> 3677

<212> DNA

<213> Artificial Sequence

<220>

<223> derived from *Arabidopsis thaliana*

<400> 15

aattattcac tcaaattcac aaagggtttc gactttgctc cgaggaagaa gataatatga 60

aaagagcttt	ttaggggttta	tcattctcct	tgactttgca	aaacgtgaaa	taagcttctt	120
ccttggagga	ctaatcaaga	cagaaatctg	ttcctctaaa	aacgatcgcc	gttctagtgc	180
actcaaatga	tgggtgaaagt	tacaaagctt	gtggcttcac	gtccaattgt	ggtcttttgc	240
gtcctggcat	tcctgggtgg	tgttttcgag	tgcatTTTgga	tctcaaattg	gcgaacaaca	300
acggagaacc	tagtcaaaga	ggtcgcttca	tttaccgaag	atctccggac	aagtctagtt	360
tcggagattg	aaaacatcgg	aaaattttaca	tatgctaaga	caaacttatc	tacgatcgg	420
ttagcgagag	ttatagattc	ttatatcacc	aacaacgaca	ctggtttttac	agagattcaa	480
acacagatcg	caccattggt	gtttgtagct	tattcaacga	tccttcaagt	ctcacaagtt	540
tcgtacatca	gtagggacgg	tctcatgttt	tcttacattg	cagaatcaaa	cacaagtgtc	600
gctgtttttg	ccaattcctc	gtcgaattca	agtcgtggag	actacacttg	gtacactcaa	660
accgtggatc	agttaactgg	tcgtcttaac	gggaactcaa	cgaaatctca	gtcgttagat	720
gtaacccata	cagattgggt	ccaagcagca	cagagtaata	actacactac	agcctttgta	780
ggaacgagct	tgggaggaga	agataacgag	actctaatac	agagcgtgg	tagcttgtac	840
agcaagaaag	gtcttggttt	tttaggggtt	ccggtttaaga	ctttaaccga	agttttgaac	900
agtttgaatc	tacacggcga	agagctttac	atgtggacaa	aggacgggac	ggtgcttgtt	960
cgtgaagggt	cactgaatga	ttctttcttc	atctccaatg	gctcgatttg	cttcggtaga	1020
gaatcgaact	ccctctgggt	tcaatgcac	cctgaaaatt	gcagttccag	tggctacgag	1080
gtggagatca	aaagattaag	ataccaagct	ttttgctctg	ttattgaagt	ttcgggctgt	1140
cctctgagat	acacactcat	gtttcccaac	aaaggaggag	caacacgcat	caagcaccaa	1200
gcggaaaagg	caaaatatca	acttattgtg	gttatgatat	ttcttggctt	cggttggcct	1260
gtatggtttg	tgtggtttat	gatgcaagca	acaaggagag	agatgcatat	gcgtgcaacg	1320
ctgataaacc	aatggaagc	gacacaacaa	gctgagagaa	agagcatgaa	caagagtcaa	1380
gcatttgcaa	atgctagcca	cgatattaga	ggtgcccttg	cagggatgaa	aggtctgatt	1440
gatatatgtc	gtgatggagt	taaacctggc	tccgacgtag	acaccactct	caaccaagtg	1500
aatgtttgcg	ccaaggattt	ggttgctctg	ctcaactctg	ttttggacat	gagcaaaatc	1560
gaaagcggga	agatgcagtt	agtggaaaga	gatttcaact	tgtcgaaact	tcttgaagac	1620
gtcatcgatt	tttaccatcc	cgttgcgatg	aagaaagggg	ttgatgtagt	tttggatccg	1680
cacgatggct	cggttttcaa	attctcgaat	gtacgagggg	atagtggcag	actgaagcag	1740
attctgaaca	atcttggttag	caatgctgtc	aagttcaccc	tcgacgggca	cattgcggta	1800
agagcttggtg	ctcagaggcc	aggttccaat	agctctgtgg	tccttgcac	atatacctaaa	1860
ggtgtgtcca	agtttgtaaa	gagtatgttc	tgcaagaata	aagaagagtc	atcaacctac	1920
gagacagaaa	tatcgaattc	cataagaaac	aatgcaaaca	cgatggagtt	tgtgtttgaa	1980
gtggatgata	ctggtaaaag	gatacctatg	gagatgcgta	agtcgggtatt	tgaaaactat	2040
gttcaagtaa	gagaaacagc	tcaaggacac	caaggaactg	gttttagggct	cgggattgtg	2100
cagtcttttg	taagattaat	gggagggggg	ataagaatca	ccgacaaggc	catgggagag	2160
aaaggaacat	gtttccaatt	caatgtttta	ttgacaacat	tagagtctcc	tccagtgagt	2220
gacatgaaag	tgagacagga	gatcgaagca	ggaggcgatt	atgtatccac	gccaaacctc	2280
gggctgacta	taaacacttc	acttgagggt	agcatgaata	tacgtaacct	gagtcctaga	2340
ttcaacaact	gtctcagctc	aagtccaaag	caagaagggt	ctagagtgg	tcttctattg	2400
aaaaatgaag	aacgcagaag	agttacagag	aaatacataa	agaatcttgg	gattaaagtt	2460
acagtgggtg	agaaatggga	gcatttgagt	tatgcttttg	agagactttt	tggattttca	2520
cctcagagtt	ccatgggaag	agcagagtgt	agtttgtcat	gtccgagctc	aagggagtta	2580
cctttcattg	gcatggacgg	tattgattca	agaagccaac	ttcctaaaag	gagaagcatc	2640
agtttctctg	cagttgtcct	tttggtgatt	gatgcaaaaa	ctggaocatt	ttttgagctg	2700
tgcgatattg	tcaaacagtt	tcgtagagge	ttgccccatg	gaatatcctg	taaagttggt	2760
tggcttaacg	aatcgagcac	tcgtgtaagt	gagagagggg	acattagttg	ttcgagacct	2820
ttgcacggat	cgcgtcttat	ggaagtcttg	aagatgttgc	ctgaatttgg	aggaaactgtg	2880
ctaaaagaac	cacctactga	gctgcaaagg	gaatcattgc	tgagacattc	ttttgttgca	2940
gagagatcac	caaaacataa	agtccaagaa	gaggggcca	gctcaatgtt	taacaaaaaa	3000
ttaggttaaga	ggataatggc	atcaacagat	tcagagagtg	agactaggg	caagtcagtg	3060
cgtacgggtc	gaaagcctat	tgggaaccca	gaggacgagc	aagagacttc	caagccgagt	3120
gacgatgaat	tcttaagagg	aaagagagtt	cttgtggctg	atgataactt	tatatcacgt	3180
aaagttgcaa	caggaaagct	gaagaagatg	ggagtctcag	aggtcgaaca	atgcgacagt	3240
gggaaagaag	ctttgagatt	agtcactgaa	gggcttacac	aaagagaaga	acaaggttca	3300
gtagataaac	ttccgtttga	ctacatatc	atgaactgcc	aatgccaga	aatggatggc	3360
tatgaagcaa	ctagagagat	taggaaagtg	gagaaaagtt	atgggggtgcg	tacaccaatt	3420
atagctgtat	ctggtcatga	tcctgggttca	gaggaagcaa	gagaaacat	tcaagctgga	3480
atggacgcct	tcttagataa	aagcttgaat	caacttgcaa	acgtcattag	agaaatcgaa	3540

```

agcaaacgtc actagtattg tataagtaga cagcacgcag ctttgtgcta ctaccattag 3600
tgtttttagac aatacttttt tagtatgacg atgtcatagc agttttcttc taagtttatt 3660
taaagtccat ttcagat                                     3677

```

<210> 16

<211> 3531

<212> DNA

<213> *Arabidopsis thaliana*

<400> 16

```

atgtctataa cttgtgagct cttgaatctt acttcaaaga aagctaagaa gtcgtcgagc 60
agtgaacaaga aatggctaaa gaagcctctc ttcttctcga ttttgtgtgg ctctttggta 120
attgttttgg ttatgttctt acggttaggt agaagtcaga aggaggagac agattcttgt 180
aatggagaag agaaagtgtt gtatagacat caaaatgtca caagaagtga gattcatgat 240
ttggtctctt tgttctctga ttcagatcag gtaacatcct ttgaatgtca taaggaatca 300
agccctggaa tgtggacaaa ctatggtatt acatgttccc tgagtgtgcg ttctgataaa 360
caagagacta gagggttcc ctggaatctt ggcttaggac attctatctc atcaacatct 420
tgtatgtgtg gtaatcttga accgatttta cagcaacctg aaaaccttga ggaagaaaac 480
catgaagaag ggctggagca gggtttgtca tcgtatttaa gaaatgcatg gtggtgtcta 540
atccttgggtg tgttagtgtg ccataagatt tatgtatctc attctaaagc acgaggtgag 600
aggaaagaga aagtacatct gcaagaggct ttagctccaa agaagcagca acaacgtgct 660
cagacttctt ctagaggggc tggaagatgg aggaagaata tccttctcct tggattttta 720
ggaggagtgt ccttctctgt ttggtgggtt tgggacacta atgaggagat cataatgaaa 780
aggagggaga ctttggcaaa catgtgtgac gaacgagcac gtgttttaca agatcagttc 840
aatgttagct tgaacctatg tcatgccttg tctattcttg tatctacatt tcatcatggt 900
aaaatcccat ctgccattga tcagagaaca tttgaagaat atactgagag aacaaacttt 960
gagaggccac ttactagtgg tgtagcgtat gctttgaaag tcccacactc agaaagagag 1020
aaatttgaaa aggagcatgg atgggcaata aagaaaatgg aaactgagga ccagacagtt 1080
gtacaagatt gtgttcttga aaactttgat cccgcaccga ttcaagacga atacgcgcca 1140
gttatatttg ctcaagaaac tgtttcccat attgtatcgg tcgacatgat gtctggagaa 1200
gaagaccgtg aaaacatctt acgggcaagg gcatcaggaa aaggagtgtt aacatctcca 1260
tttaagcttc ttaagtcaaa tcatcttggg gttgtgttga cctttgctgt ctatgacacg 1320
agcctaccgc ctgatgctac agaagaacag cgtgttgaag caactattgg gtaccttggg 1380
gcatcatatg atatgccatc gctgggtggag aaacttcttc accaacttgc cagcaaacag 1440
acaattgctg tggatgttta cgacacaact aacacttcag gtctaataaa aatgtatggc 1500
tcagaaattg gggatataag tgagcagcat ataagtagcc ttgatttttg tgatccatca 1560
aggaacctat agatgcattg caggtttaag cataaacttc ccattccctg gacagcgata 1620
acaccgtcga tcttagttct ggttattact tttcttgttg gttatatatt atatgaagcc 1680
atcaaccgaa ttgcgacagt tgaagaggat tgtcagaaga tgagggaact caaagctcgt 1740
gctgaggccg ctgacattgc aaagtcacag ttcctagcaa ctgtttctca tgagatacgg 1800
actccgatga atggagtgtt aggaatgctg aaaatgctga tggacaccga tcttgatgctg 1860
aagcagatgg actatgcgca aactgctcat ggcagtggga aggatcttac atcactaata 1920
aatgaggttc ttgatcaggc aaagattgaa tccggaaggc tcgagcttga aaatgtgcct 1980
tttgatatgc gttttattct tgataatgtt tcatctctcc tctctggcaa ggcaaataaa 2040
aaaggaattg agttggccgt ttatgtttct agtcaagttc ctgatgttgt agtcggtgat 2100
ccgagtcggt tccggcagat cattacaaac ctggttggaa actcaatcaa attcacacag 2160
gaaaggggac acatatttat ctcatgtcac cttgcagatg aggtaaagga gcctcttact 2220
attgaagacg cagtgtctaaa acagcgacta gctttaggat gcagcgagtc cggtgagaca 2280
gttagcgggt ttcttgcggt aaatgcatgg ggaagctgga agaatttcaa gacatgttac 2340
agtactgaga gtcagaattc tgatcaaate aaattgctag ttacagtgga ggacactgga 2400
gttggcatat ctgtggatgc acaaggccga atcttcacac cttttatgca agccgacagt 2460
tccacatcgc ggacttatgg tggaactggc ataggtttga gtataagcaa acgtttgggt 2520
gaactcatgc aaggagagat ggggtttgtg agtgagcccg ggataggcag tactttttca 2580
tttactggag ttttcgggaa agcagaaaca aatacgtcga ttactaagct ggaacgattc 2640
gatctagcta ttcaggagtt tacaggattg agagcattag ttattgataa cagaaacatt 2700
agagcagagg tcaccaggta cgaacttcgg agactgggaa tatctgcaga cattgtttca 2760
agtctgagaa tggcatgcac ttgttgtatc agcaaattag aaaatttggc tatgattcta 2820
atagacaaag acgcctggaa caaggaagaa ttttcagtag ttgacgagtt gtttaccgca 2880

```

```

agcaaagtaa cctttacaag agtcccaaag atttttcttt tggcaacttc tgcaactctt 2940
actgagcgca gtgagatgaa gtctactggg ctcactgatg aggtgggtgat aaagcctctt 3000
cggatgagtg tcttaatatg ttgcttgcaa gaaacccttg tcaatggcaa gaagaggcaa 3060
ccgaacagac agcgaagaaa tcttgggacac ttgctaagag aaaaacagat tctggttgtg 3120
gatgataatc ttgtgaacag acgagttgca gaaggtgcac ttaagaaata tggagctatt 3180
gttacatgcg ttgagagtgg caaagctgca ttggcaatgc ttaagccgcc tcataacttc 3240
gatgcttgct tcatggatct ccagatgcct gaaatggatg gatttgaagc gacaaggaga 3300
gtccgtgagc tggagaggga aatcaataag aaaatagctt ctggagaagt ttcagctgaa 3360
atgttctgta aatttagtag ttggcacgtc ccgatattag caatgacagc agatgttatt 3420
caggctactc atgaagaatg catgaaatgt ggaatggatg gttatgtatc aaaaccgttt 3480
gaagaggaag tgctctacac agcggtagca agattctttg aaccttggtta a 3531

```

<210> 17

<211> 1503

<212> DNA

<213> *Arabidopsis thaliana*

<400> 17

```

atggctccag taataaagct ggtcttaggt tcagttgcct ttgccatttt ctggatatta 60
gcagtgtttc cctctgtacc attcctacca attggctcgaa ccgccgggtc tctgttcggg 120
gcaatgctaa tgggttatctt ccaagtgata accccggaac aagcttatgc agccattgat 180
ctcccagatac tcggtcttct ctttgggaacg atgggtgtta gtatatacct tgagagagct 240
gatatgttca agtacttagg cacattgctt tcgtggaaaa gcagagggtc gaaagacttg 300
ctatgtagag tctgtcttgt ttcegtgtt tcaagcgctc tgtttactaa cgatacatct 360
tgtgtggttt taaccgagtt tgtgttgaag atcgctaggc aaaagaatct cccgcctcac 420
ccgtttttgc tcgcttttagc cacgagtgcg aatattgggt cttctgctac tccattggg 480
aatccgcaga atcttgttat tgccgttcaa agcaagatcc cgttttgagg gtttcttctt 540
gggtgttttc ctgcgatgat tgttggtatt accgttaacg ctatgcttct tctcggtatg 600
tattggcggg tattgtcaga tcataaagag gatgaagaag aagtacaaaa tgctgattct 660
gaagttgttg cccaagaaga tgtccaatct cataggttct cgccagctac attttcgcca 720
gtctcatccg aagattcgaa cttgagaatg gatgccgccg agactctcag aaacagagcg 780
ggttcagcgg gtgagagcga gttaataagc tgcaattcaa atgcgtcgag agaacaacac 840
aacgacgcag agtctcaagg agagagcaat aataccaaca atatgttcca gaccaagaga 900
tggaagaagag ttttatggaa atcaagtgtt tatttcatca cattaggaat gctaatatct 960
ctgcttatgg gtttaaacat gtcgtggacc gcaattaccg cggctctagc actcgtgggt 1020
cttgatttca aagacgcaag gccgtctctc gagaagggtat cgtattcgct tttgatcttt 1080
ttctgcggga tgttcataac cgttgatgga ttcaacaaaa ctggtatccc taccgctcta 1140
tgggacctaa tggagccata tgccaaaatt gatcaagcca aaggaatcgc ggttctagca 1200
gttgtgattc ttgtcctctc caatgtagcc tctaattgtac caaccgtgct tttgttgagg 1260
gcaagagtgg cggcatcagc gatggggagg gaggaggaga agaaggcgtg gttgttgctg 1320
gcgtgggtga gcacagtggc tggaaacttg acgttgcttg gttcagcagc aaacctgata 1380
gtgtgtgagc aagctcgtag agcagtgagc catggataca ctctcacttt cactaaacat 1440
ttcaaatcgc gtttgccttc aacactcatc gtcactgcga ttgggtctct ccttatcaag 1500
taa 1503

```

<210> 18

<211> 1803

<212> DNA

<213> Artificial Sequence

<220>

<223> derived from *Arabidopsis thaliana*

<400> 18

```

atgcacatag taaaagtcga agatgatttc catgaaatgc aagagcttaa agttcgagca 60
gaagctgctg atgtcgctaa atcgcagttt cttgctaccg tgtctcacga gatcaggaca 120
ccaatgaatg gcattctcgg aatgcttgct atgctcctag atacagaact aagctcgaca 180
cagagagatt acgctcaaac cgctcaagta tgtggtaaag ctttgattgc attgataaat 240

```

```

gaggttcttg atcgcgccaa gattgaagct ggaaagctgg agttggaatc agtaccattt 300
gatatccgtt caatattgga tgatgtcctt tctctattct ctgaggagtc aaggaacaaa 360
agcattgagc tcgcgggttt cgtttcagac aaagtaccag agatagtcaa aggagattca 420
gggagattta gacagataat cataaacctt gttggaaatt cggttaaatt cacagagaaa 480
ggacatatct ttgttaaagt ccatcttgcg gaacaatcaa aagatgaatc tgaaccgaaa 540
aatgcattga atggtggagt caatgttagg cttatgggtt caatcgaaga cacgggtatt 600
ggaatccctt tagttgcgca aggccgtgtg tttatgccgt ttatgcaagc agatagctcg 660
acttcaagaa actatggagg tactggtatt ggtttgagta taagcaagtg tcttggtgaa 720
cttatgcgtg gtcagataaa tttcataagc cggcctcata ttggaagcac gttctgggtc 780
acggctgttt tagagaaatg cgataaatgc agtgcgatta accatatgaa gaaacctaat 840
gtggaacact tgccttctac ttttaaagga atgaaagcta tagttgttga tgctaagcct 900
gttagagctg ctgtgactag ataccatatg aaaagactcg gaatcaatgt tgatgtcgtg 960
acaagtctca aaaccgctgt tgttgcagct gctgcgtttg aaagaaacgg ttctcctctc 1020
ccaacaaaac cgcaacttga tatgatctta gtagagaaag attcatggat ttcaactgaa 1080
gataatgact cagagattcg tttattgaat tcaagaacca acggaaacgt tcatcacaag 1140
tctccgaaac tagctctatt cgcaacaaac atcacaaatt cggagttcga cagagctaaa 1200
tccgcaggat ttgcagatac ggtaataatg aaaccgttaa gagcaagcat gattggggcg 1260
tgtctgcaac aagttctcga gctgagaaaa acaagacaac aacatccaga aggatcatca 1320
cccgcactc tcaagagctt gcttacaggg aagaagattc ttgtggttga tgataatata 1380
gttaacagga gagtagctgc aggagctctc aagaaatttg gagcagaagt ggtttgtgca 1440
gagagtggtc aagttgcttt gggtttgctt cagattccac acactttcga tgcttgcttc 1500
atggatattc aaatgccaca gatggacgga atgatggaga aggaaactaa agagaagaca 1560
aatctcgaat ggcatttacc gattctagcg atgactgcgg atgtgataca cgcgacctac 1620
gaggaatgtc tgaaaagtgg gatggatggg tacgtctcca aaccttttga agaagagaat 1680
ctctataaat ccgttgccaa atcattcaaa cctaatacta tctcaccttc ccggaagagt 1740
ggaccgatct tgctgattga tatgcatttt ggtttctgta catacagtag gttcacaatc 1800
tag 1803

```

<210> 19

<211> 1803

<212> DNA

<213> Artificial Sequence

<220>

<223> derived from *Arabidopsis thaliana*

<400> 19

```

atgcacatag taaaagtcga agatgatttc catgaaatgc aagagcttaa agttcgagca 60
gaagctgctg atgtcgctaa atcgcagttt cttgctaccg tgtctcaaga gatcaggaca 120
ccaatgaatg gcattctcgg aatgcttgct atgctcctag atacagaact aagctcgaca 180
cagagagatt acgctcaaac cgctcaagta tgtggtaaag ctttgattgc attgataaat 240
gaggttcttg atcgcgccaa gattgaagct ggaaagctgg agttggaatc agtaccattt 300
gatatccgtt caatattgga tgatgtcctt tctctattct ctgaggagtc aaggaacaaa 360
agcattgagc tcgcgggttt cgtttcagac aaagtaccag agatagtcaa aggagattca 420
gggagattta gacagataat cataaacctt gttggaaatt cggttaaatt cacagagaaa 480
ggacatatct ttgttaaagt ccatcttgcg gaacaatcaa aagatgaatc tgaaccgaaa 540
aatgcattga atggtggagt caatgttagg cttatgggtt caatcgaaga cacgggtatt 600
ggaatccctt tagttgcgca aggccgtgtg tttatgccgt ttatgcaagc agatagctcg 660
acttcaagaa actatggagg tactggtatt ggtttgagta taagcaagtg tcttggtgaa 720
cttatgcgtg gtcagataaa tttcataagc cggcctcata ttggaagcac gttctgggtc 780
acggctgttt tagagaaatg cgataaatgc agtgcgatta accatatgaa gaaacctaat 840
gtggaacact tgccttctac ttttaaagga atgaaagcta tagttgttga tgctaagcct 900
gttagagctg ctgtgactag ataccatatg aaaagactcg gaatcaatgt tgatgtcgtg 960
acaagtctca aaaccgctgt tgttgcagct gctgcgtttg aaagaaacgg ttctcctctc 1020
ccaacaaaac cgcaacttga tatgatctta gtagagaaag attcatggat ttcaactgaa 1080
gataatgact cagagattcg tttattgaat tcaagaacca acggaaacgt tcatcacaag 1140
tctccgaaac tagctctatt cgcaacaaac atcacaaatt cggagttcga cagagctaaa 1200
tccgcaggat ttgcagatac ggtaataatg aaaccgttaa gagcaagcat gattggggcg 1260

```



```

tgtctgcaac aagttctcga gctgagaaaa acaagacaac aacatccaga aggatcatca 1320
cccgcaactc tcaagagctt gcttacaggg aagaagattc ttgtgggttga tgataatata 1380
gttaacagga gagtagctgc aggagctctc aagaaatttg gagcagaagt ggtttgtgca 1440
gagagtggtc aagttgcttt gggtttgctt cagattccac acactttcga tgcttgcttc 1500
atggatattc aaatgccaca gatggacgga atgatggaga aggaaactaa agagaagaca 1560
aatctcgaat ggcattttacc gattctagcg atgactgceg atgtgatata cgcgacctac 1620
gaggaatgtc tgaaaagtgg gatggatggg tacgtctcca aaccttttga agaagagaat 1680
ctctataaat ccgttgccaa atcattcaaa cctaatacta tctcaccttc ccggaagagt 1740
ggaccgattc tgctgattga tatgcatttt ggtttctgta catacagtag gttcacaatc 1800
tag

```

<210> 20

<211> 1803

<212> DNA

<213> Artificial Sequence

<220>

<223> derived from *Arabidopsis thaliana*

<400> 20

```

atgcacatag taaaagtcga agatgatttc catgaaatgc aagagcttaa agttcgagca 60
gaagctgctg atgtcgctaa atcgagcttt cttgctaccg tgtctcacga gatcaggaca 120
ccaatgaatg gcattctcgg aatgcttgct atgctcctag atacagaact aagctcgaca 180
cagagagatt acgctcaaac cgctcaagta tgtggtaaag ctttgattgc attgataaat 240
gaggttcttg atcgcgccaa gattgaagct ggaaagctgg agttggaatc agtaccattt 300
gatatccgtt caatattgga tgatgtcctt tctctattct ctgaggagtc aaggaacaaa 360
agcattgagc tcgcggtttt cgtttcagac aaagtaccag agatagtcaa aggagattca 420
gggagattta gacagataat cataaacctt gttggaaatt cgggttaaatt cacagagaaa 480
ggacatatct ttgttaaagt ccatcttgcg gaacaatcaa aagatgaatc tgaaccgaaa 540
aatgcattga atgggtggagt caatgttagg cttatgggtt caatcgaaga cacgggtatt 600
ggaatccctt tagttgcgca aggcctgttg tttatgccgt ttatgcaagc agatagctcg 660
acttcaagaa actatggagg tactgggtatt ggtttgagta taagcaagtg tcttggtgaa 720
cttatgcgtg gtcagataaa tttcataagc cggcctcata ttggaagcac gttctggttc 780
acggctggtt tagagaaatg cgataaatgc agtgcgatta accatatgaa gaaacctaat 840
gtggaacact tgccttctac ttttaaagga atgaaagcta tagttgttga tgctaagcct 900
gttagagctg ctgtgactag ataccatatg aaaagactcg gaatcaatgt tgatgtcgtg 960
acaagtctca aaaccgctgt tgttgagct gctgcgtttg aaagaaacgg ttctcctctc 1020
ccaacaaaac cgcaacttga tatgatctta gtagagaaag attcatggat ttcaactgaa 1080
gataatgact cagagattcg tttattgaat tcaagaacca acggaaacgt tcatcacaag 1140
tctccgaaac tagctctatt cgcaacaaac atcacaaatt cggagttcga cagagctaaa 1200
tccgcaggat ttgcagatac ggtaataatg aaacogttaa gagcaagcat gattggggcg 1260
tgtctgcaac aagttctcga gctgagaaaa acaagacaac aacatccaga aggatcatca 1320
cccgcaactc tcaagagctt gcttacaggg aagaagattc ttgtgggttga tgataatata 1380
gttaacagga gagtagctgc aggagctctc aagaaatttg gagcagaagt ggtttgtgca 1440
gagagtggtc aagttgcttt gggtttgctt cagattccac acactttcga tgcttgcttc 1500
atgaatattc aaatgccaca gatggacgga atgatggaga aggaaactaa agagaagaca 1560
aatctcgaat ggcattttacc gattctagcg atgactgceg atgtgatata cgcgacctac 1620
gaggaatgtc tgaaaagtgg gatggatggg tacgtctcca aaccttttga agaagagaat 1680
ctctataaat ccgttgccaa atcattcaaa cctaatacta tctcaccttc ccggaagagt 1740
ggaccgattc tgctgattga tatgcatttt ggtttctgta catacagtag gttcacaatc 1800
tag

```

<210> 21

<211> 465

<212> DNA

<213> *Arabidopsis thaliana*

<400> 21

```

atggatttgg ttcagaagca gaagagtttg caagattaca ccaaactcact cttcttagaa 60
gggatttgg acagccagtt cttgcagctg caacaactac aagatgaaag caatccagat 120
tttgtttcac aagttgtcac actcttcttc caagactctg ataggattct caatgatctc 180
tcactttccc tagatcaaca agttgtagac tttaaaaaag ttgatcccca tgttcatcaa 240
ctcaaaggta gcagctccag tataggagca cagagagtta agaatgcttg tgttgtcttc 300
cgcagcttct gcgagcagca aaatgtcgaa gcatgtcata gatgtttgca acaagtgaag 360
caagagtatt atcttgtgaa aaacagatta gagactctgt tcaagctgga gcaacagatt 420
gtagcttctg gtggaatgat cccggccgctc gaactcggat tttga 465

```

<210> 22

<211> 817

<212> DNA

<213> *Arabidopsis thaliana*

<400> 22

```

gaaagacaaa acacaagttt cttcttcttg ggattggcta tttccagaaa tccaagtcaa 60
taatcaaagt ccaaacaaaa aaatcctctc ccaatctccg cttcactctt ctcatggacg 120
ctctcattgc tcagcttcag agacaatttc gtgattacac catttctctc taccaacagg 180
ggtttttggg tgatcaattt actgagttga aaaagctaca agatgatgga agtcctgatt 240
ttgtgtctga agtgctttca ctttctcttg aagatttgtt gaagcttatc agtaacatgg 300
ctagagcttt ggacacgaca ggaactgtag attttagtca ggtaggtgct agtgtgcatc 360
aattgaaggg tagtagctca agtggttggtg ccaagagggg caaaactttg tgtgttagct 420
tcaaggaatg ttgtgaagct aagaactacg aagggtgtgt gagatgtttg cagcaagtgg 480
atattgagta caaggcgtta aagacaaagc ttcaagatat gttcaatctt gagaaacaga 540
tcattcaagc tgggtgtata gtccctcaag tggatattaa cttaaagagac tagtccataa 600
gaagaaaaaa gatgatgact ttcttctctt agtttctctt cttaaattatt ttggatttgg 660
tgtttgctca aaaactcaat aaaatatgtg caaaaagaaa caaaaacaag tgatggttgt 720
ttataaatca gtagtatgta ttgtttgatc tcatccgaga aaattgaaac cattggacta 780
atgaatgtga tgataatata tattggtttg cttctcc 817

```

<210> 23

<211> 510

<212> DNA

<213> *Arabidopsis thaliana*

<400> 23

```

aacttttagct atgaacacca tcgtcgttgc tcagttgcag agacaatttc aagactacat 60
cgttttctctt tatcaacagg gatttctgga taatcagttc tcagagttga gaaagttgca 120
agatgaagga acccctgatt ttgtagctga agttgtctct ctattcttcg acgactgttc 180
caagcttatt aataccatgt ctatatccct ggagcggcca gataatgtgg atttcaaaca 240
ggtggattca ggtgttcac aactcaaggg tagtagctcc agtgctcggg caaggagggg 300
gaaaaatgtg tgcatactt tcaaggaatg ttgcgatgtt cagaaccgtg aagggtgtct 360
aagggtgttta cagcaggtgg attatgaata taagatgtta aagactaaac ttcaggatct 420
ctttaattta gagaaacaga tcctccaagc tggaggtaca attcctcaag tggatataaa 480
ttagaccgat gcgtttctcg attatgcaaa 510

```

<210> 24

<211> 465

<212> DNA

<213> Artificial Sequence

<220>

<223> derived from *Arabidopsis thaliana*

<400> 24

```

atggatttgg ttcagaagca gaagagtttg caagattaca ccaaactcact cttcttagaa 60
gggatttgg acagccagtt cttgcagctg caacaactac aagatgaaag caatccagat 120
tttgtttcac aagttgtcac actcttcttc caagactctg ataggattct caatgatctc 180

```

```

tcactttccc tagatcaaca agttgtagac tttaaaaaag ttgatcccca tgttcaacaa 240
ctcaaaggta gcagctccag tataggagca cagagagtta agaatgcttg tgttgtcttc 300
cgcagcttct gcgagcagca aaatgtcgaa gcatgtcata gatgtttgca acaagtgaag 360
caagagtatt atcttgtgaa aaacagatta gagactctgt tcaagctgga gcaacagatt 420
gtagcttctg gtggaatgat cccggccgctc gaactcggat tttga 465

```

<210> 25

<211> 465

<212> DNA

<213> Artificial Sequence

<220>

<223> derived from *Arabidopsis thaliana*

<400> 25

```

atggatttgg ttcagaagca gaagagtttg caagattaca ccaaactact cttcttagaa 60
gggatttgg acagccagtt cttgcagctg caacaactac aagatgaaag caatccagat 120
tttgtttcac aagttgtcac actcttcttc caagactctg ataggattct caatgatctc 180
tcactttccc tagatcaaca agttgtagac tttaaaaaag ttgatcccca agttcaacaa 240
ctcaaaggta gcagctccag tataggagca cagagagtta agaatgcttg tgttgtcttc 300
cgcagcttct gcgagcagca aaatgtcgaa gcatgtcata gatgtttgca acaagtgaag 360
caagagtatt atcttgtgaa aaacagatta gagactctgt tcaagctgga gcaacagatt 420
gtagcttctg gtggaatgat cccggccgctc gaactcggat tttga 465

```

<210> 26

<211> 1176

<212> PRT

<213> *Arabidopsis thaliana*

<400> 26

```

Met Ser Ile Thr Cys Glu Leu Leu Asn Leu Thr Ser Lys Lys Ala Lys
 1          5          10          15
Lys Ser Ser Ser Ser Asp Lys Lys Trp Leu Lys Lys Pro Leu Phe Phe
          20          25          30
Leu Ile Leu Cys Gly Ser Leu Val Ile Val Leu Val Met Phe Leu Arg
          35          40          45
Leu Gly Arg Ser Gln Lys Glu Glu Thr Asp Ser Cys Asn Gly Glu Glu
          50          55          60
Lys Val Leu Tyr Arg His Gln Asn Val Thr Arg Ser Glu Ile His Asp
65          70          75          80
Leu Val Ser Leu Phe Ser Asp Ser Asp Gln Val Thr Ser Phe Glu Cys
          85          90          95
His Lys Glu Ser Ser Pro Gly Met Trp Thr Asn Tyr Gly Ile Thr Cys
          100          105          110
Ser Leu Ser Val Arg Ser Asp Lys Gln Glu Thr Arg Gly Leu Pro Trp
          115          120          125
Asn Leu Gly Leu Gly His Ser Ile Ser Ser Thr Ser Cys Met Cys Gly
          130          135          140
Asn Leu Glu Pro Ile Leu Gln Gln Pro Glu Asn Leu Glu Glu Glu Asn
145          150          155          160
His Glu Glu Gly Leu Glu Gln Gly Leu Ser Ser Tyr Leu Arg Asn Ala
          165          170          175
Trp Trp Cys Leu Ile Leu Gly Val Leu Val Cys His Lys Ile Tyr Val
          180          185          190
Ser His Ser Lys Ala Arg Gly Glu Arg Lys Glu Lys Val His Leu Gln
          195          200          205
Glu Ala Leu Ala Pro Lys Lys Gln Gln Gln Arg Ala Gln Thr Ser Ser
          210          215          220

```


Arg	Gly	Ala	Gly	Arg	Trp	Arg	Lys	Asn	Ile	Leu	Leu	Leu	Gly	Ile	Leu
225					230					235					240
Gly	Gly	Val	Ser	Phe	Ser	Val	Trp	Trp	Phe	Trp	Asp	Thr	Asn	Glu	Glu
				245					250					255	
Ile	Ile	Met	Lys	Arg	Arg	Glu	Thr	Leu	Ala	Asn	Met	Cys	Asp	Glu	Arg
			260					265					270		
Ala	Arg	Val	Leu	Gln	Asp	Gln	Phe	Asn	Val	Ser	Leu	Asn	His	Val	His
		275					280					285			
Ala	Leu	Ser	Ile	Leu	Val	Ser	Thr	Phe	His	His	Gly	Lys	Ile	Pro	Ser
	290					295					300				
Ala	Ile	Asp	Gln	Arg	Thr	Phe	Glu	Glu	Tyr	Thr	Glu	Arg	Thr	Asn	Phe
305					310				315						320
Glu	Arg	Pro	Leu	Thr	Ser	Gly	Val	Ala	Tyr	Ala	Leu	Lys	Val	Pro	His
				325					330					335	
Ser	Glu	Arg	Glu	Lys	Phe	Glu	Lys	Glu	His	Gly	Trp	Ala	Ile	Lys	Lys
			340					345					350		
Met	Glu	Thr	Glu	Asp	Gln	Thr	Val	Val	Gln	Asp	Cys	Val	Pro	Glu	Asn
		355					360					365			
Phe	Asp	Pro	Ala	Pro	Ile	Gln	Asp	Glu	Tyr	Ala	Pro	Val	Ile	Phe	Ala
	370					375					380				
Gln	Glu	Thr	Val	Ser	His	Ile	Val	Ser	Val	Asp	Met	Met	Ser	Gly	Glu
385					390					395					400
Glu	Asp	Arg	Glu	Asn	Ile	Leu	Arg	Ala	Arg	Ala	Ser	Gly	Lys	Gly	Val
				405					410					415	
Leu	Thr	Ser	Pro	Phe	Lys	Leu	Leu	Lys	Ser	Asn	His	Leu	Gly	Val	Val
			420					425					430		
Leu	Thr	Phe	Ala	Val	Tyr	Asp	Thr	Ser	Leu	Pro	Pro	Asp	Ala	Thr	Glu
		435					440					445			
Glu	Gln	Arg	Val	Glu	Ala	Thr	Ile	Gly	Tyr	Leu	Gly	Ala	Ser	Tyr	Asp
	450					455					460				
Met	Pro	Ser	Leu	Val	Glu	Lys	Leu	Leu	His	Gln	Leu	Ala	Ser	Lys	Gln
465					470					475					480
Thr	Ile	Ala	Val	Asp	Val	Tyr	Asp	Thr	Thr	Asn	Thr	Ser	Gly	Leu	Ile
				485					490					495	
Lys	Met	Tyr	Gly	Ser	Glu	Ile	Gly	Asp	Ile	Ser	Glu	Gln	His	Ile	Ser
		500						505					510		
Ser	Leu	Asp	Phe	Gly	Asp	Pro	Ser	Arg	Asn	His	Glu	Met	His	Cys	Arg
		515					520					525			
Phe	Lys	His	Lys	Leu	Pro	Ile	Pro	Trp	Thr	Ala	Ile	Thr	Pro	Ser	Ile
	530					535					540				
Leu	Val	Leu	Val	Ile	Thr	Phe	Leu	Val	Gly	Tyr	Ile	Leu	Tyr	Glu	Ala
545					550					555					560
Ile	Asn	Arg	Ile	Ala	Thr	Val	Glu	Glu	Asp	Cys	Gln	Lys	Met	Arg	Glu
				565					570					575	
Leu	Lys	Ala	Arg	Ala	Glu	Ala	Ala	Asp	Ile	Ala	Lys	Ser	Gln	Phe	Leu
		580						585					590		
Ala	Thr	Val	Ser	His	Glu	Ile	Arg	Thr	Pro	Met	Asn	Gly	Val	Leu	Gly
		595					600					605			
Met	Leu	Lys	Met	Leu	Met	Asp	Thr	Asp	Leu	Asp	Ala	Lys	Gln	Met	Asp
	610					615					620				
Tyr	Ala	Gln	Thr	Ala	His	Gly	Ser	Gly	Lys	Asp	Leu	Thr	Ser	Leu	Ile
625					630					635					640
Asn	Glu	Val	Leu	Asp	Gln	Ala	Lys	Ile	Glu	Ser	Gly	Arg	Leu	Glu	Leu
				645					650					655	
Glu	Asn	Val	Pro	Phe	Asp	Met	Arg	Phe	Ile	Leu	Asp	Asn	Val	Ser	Ser
			660					665					670		
Leu	Leu	Ser	Gly	Lys	Ala	Asn	Glu	Lys	Gly	Ile	Glu	Leu	Ala	Val	Tyr
		675					680					685			

Val	Ser	Ser	Gln	Val	Pro	Asp	Val	Val	Val	Gly	Asp	Pro	Ser	Arg	Phe
	690					695					700				
Arg	Gln	Ile	Ile	Thr	Asn	Leu	Val	Gly	Asn	Ser	Ile	Lys	Phe	Thr	Gln
705					710				715						720
Glu	Arg	Gly	His	Ile	Phe	Ile	Ser	Val	His	Leu	Ala	Asp	Glu	Val	Lys
			725						730					735	
Glu	Pro	Leu	Thr	Ile	Glu	Asp	Ala	Val	Leu	Lys	Gln	Arg	Leu	Ala	Leu
			740					745					750		
Gly	Cys	Ser	Glu	Ser	Gly	Glu	Thr	Val	Ser	Gly	Phe	Pro	Ala	Val	Asn
		755					760					765			
Ala	Trp	Gly	Ser	Trp	Lys	Asn	Phe	Lys	Thr	Cys	Tyr	Ser	Thr	Glu	Ser
770						775					780				
Gln	Asn	Ser	Asp	Gln	Ile	Lys	Leu	Leu	Val	Thr	Val	Glu	Asp	Thr	Gly
785					790					795					800
Val	Gly	Ile	Pro	Val	Asp	Ala	Gln	Gly	Arg	Ile	Phe	Thr	Pro	Phe	Met
				805					810					815	
Gln	Ala	Asp	Ser	Ser	Thr	Ser	Arg	Thr	Tyr	Gly	Gly	Thr	Gly	Ile	Gly
			820					825					830		
Leu	Ser	Ile	Ser	Lys	Arg	Leu	Val	Glu	Leu	Met	Gln	Gly	Glu	Met	Gly
		835					840					845			
Phe	Val	Ser	Glu	Pro	Gly	Ile	Gly	Ser	Thr	Phe	Ser	Phe	Thr	Gly	Val
850						855					860				
Phe	Gly	Lys	Ala	Glu	Thr	Asn	Thr	Ser	Ile	Thr	Lys	Leu	Glu	Arg	Phe
865					870					875					880
Asp	Leu	Ala	Ile	Gln	Glu	Phe	Thr	Gly	Leu	Arg	Ala	Leu	Val	Ile	Asp
				885					890					895	
Asn	Arg	Asn	Ile	Arg	Ala	Glu	Val	Thr	Arg	Tyr	Glu	Leu	Arg	Arg	Leu
			900					905					910		
Gly	Ile	Ser	Ala	Asp	Ile	Val	Ser	Ser	Leu	Arg	Met	Ala	Cys	Thr	Cys
		915					920					925			
Cys	Ile	Ser	Lys	Leu	Glu	Asn	Leu	Ala	Met	Ile	Leu	Ile	Asp	Lys	Asp
930						935					940				
Ala	Trp	Asn	Lys	Glu	Glu	Phe	Ser	Val	Leu	Asp	Glu	Leu	Phe	Thr	Arg
945					950					955					960
Ser	Lys	Val	Thr	Phe	Thr	Arg	Val	Pro	Lys	Ile	Phe	Leu	Leu	Ala	Thr
				965					970					975	
Ser	Ala	Thr	Leu	Thr	Glu	Arg	Ser	Glu	Met	Lys	Ser	Thr	Gly	Leu	Ile
			980					985					990		
Asp	Glu	Val	Val	Ile	Lys	Pro	Leu	Arg	Met	Ser	Val	Leu	Ile	Cys	Cys
		995					1000					1005			
Leu	Gln	Glu	Thr	Leu	Val	Asn	Gly	Lys	Lys	Arg	Gln	Pro	Asn	Arg	Gln
1010						1015					1020				
Arg	Arg	Asn	Leu	Gly	His	Leu	Leu	Arg	Glu	Lys	Gln	Ile	Leu	Val	Val
1025					1030					1035					1040
Asp	Asp	Asn	Leu	Val	Asn	Arg	Arg	Val	Ala	Glu	Gly	Ala	Leu	Lys	Lys
				1045					1050					1055	
Tyr	Gly	Ala	Ile	Val	Thr	Cys	Val	Glu	Ser	Gly	Lys	Ala	Ala	Leu	Ala
			1060					1065					1070		
Met	Leu	Lys	Pro	Pro	His	Asn	Phe	Asp	Ala	Cys	Phe	Met	Asp	Leu	Gln
		1075					1080					1085			
Met	Pro	Glu	Met	Asp	Gly	Phe	Glu	Ala	Thr	Arg	Arg	Val	Arg	Glu	Leu
1090						1095						1100			
Glu	Arg	Glu	Ile	Asn	Lys	Lys	Ile	Ala	Ser	Gly	Glu	Val	Ser	Ala	Glu
1105					1110					1115					1120
Met	Phe	Cys	Lys	Phe	Ser	Ser	Trp	His	Val	Pro	Ile	Leu	Ala	Met	Thr
				1125					1130					1135	
Ala	Asp	Val	Ile	Gln	Ala	Thr	His	Glu	Glu	Cys	Met	Lys	Cys	Gly	Met
			1140					1145					1150		

Asp Gly Tyr Val Ser Lys Pro Phe Glu Glu Glu Val Leu Tyr Thr Ala
 1155 1160 1165
 Val Ala Arg Phe Phe Glu Pro Cys
 1170 1175

<210> 27
 <211> 1036
 <212> PRT
 <213> Arabidopsis thaliana

<400> 27
 Met Ser Leu Phe His Val Leu Gly Phe Gly Val Lys Ile Gly His Leu
 1 5 10 15
 Phe Trp Met Leu Cys Cys Trp Phe Val Ser Trp Phe Val Asp Asn Gly
 20 25 30
 Ile Glu Asp Lys Ser Gly Leu Leu Val Gly Ser Val Gly Asp Leu Glu
 35 40 45
 Lys Thr Lys Met Thr Thr Leu Lys Lys Lys Asn Lys Met Trp Phe Trp
 50 55 60
 Asn Lys Ile Ser Ser Ser Gly Leu Lys Ile Pro Ser Phe Ser Tyr Gln
 65 70 75 80
 Phe Leu Gly Ser Val Lys Phe Asn Lys Ala Trp Trp Arg Lys Leu Val
 85 90 95
 Val Val Trp Val Val Phe Trp Val Leu Val Ser Ile Trp Thr Phe Trp
 100 105 110
 Tyr Phe Ser Ser Gln Ala Met Glu Lys Arg Lys Glu Thr Leu Ala Ser
 115 120 125
 Met Cys Asp Glu Arg Ala Arg Met Leu Gln Asp Gln Phe Asn Val Ser
 130 135 140
 Met Asn His Val Gln Ala Met Ser Ile Leu Ile Ser Thr Phe His His
 145 150 155 160
 Gly Lys Ile Pro Ser Ala Ile Asp Gln Arg Thr Phe Ser Glu Tyr Thr
 165 170 175
 Asp Arg Thr Ser Phe Glu Arg Pro Leu Thr Ser Gly Val Ala Tyr Ala
 180 185 190
 Met Arg Val Leu His Ser Glu Arg Glu Glu Phe Glu Arg Gln Gln Gly
 195 200 205
 Trp Thr Ile Arg Lys Met Tyr Ser Leu Glu Gln Asn Pro Val His Lys
 210 215 220
 Asp Asp Tyr Asp Leu Glu Ala Leu Glu Pro Ser Pro Val Gln Glu Glu
 225 230 235 240
 Tyr Ala Pro Val Ile Phe Ala Gln Asp Thr Val Ser His Val Val Ser
 245 250 255
 Leu Asp Met Leu Ser Gly Lys Glu Asp Arg Glu Asn Val Leu Arg Ala
 260 265 270
 Arg Ser Ser Gly Lys Gly Val Leu Thr Ala Pro Phe Pro Leu Ile Lys
 275 280 285
 Thr Asn Arg Leu Gly Val Ile Leu Thr Phe Ala Val Tyr Lys Arg Asp
 290 295 300
 Leu Pro Ser Asn Ala Thr Pro Lys Glu Arg Ile Glu Ala Thr Asn Gly
 305 310 315 320
 Tyr Leu Gly Gly Val Phe Asp Ile Glu Ser Leu Val Glu Asn Leu Leu
 325 330 335
 Gln Gln Leu Ala Ser Lys Gln Thr Ile Leu Val Asn Val Tyr Asp Ile
 340 345 350
 Thr Asn His Ser Gln Pro Ile Ser Met Tyr Gly Thr Asn Val Ser Ala
 355 360 365

Asp	Gly	Leu	Glu	Arg	Val	Ser	Pro	Leu	Ile	Phe	Gly	Asp	Pro	Leu	Arg
370						375					380				
Lys	His	Glu	Met	Arg	Cys	Arg	Phe	Lys	Gln	Lys	Pro	Pro	Trp	Pro	Val
385					390					395					400
Leu	Ser	Met	Val	Thr	Ser	Phe	Gly	Ile	Leu	Val	Ile	Ala	Leu	Leu	Val
				405					410						415
Ala	His	Ile	Ile	His	Ala	Thr	Val	Ser	Arg	Ile	His	Lys	Val	Glu	Glu
			420					425					430		
Asp	Cys	Asp	Lys	Met	Lys	Gln	Leu	Lys	Lys	Lys	Ala	Glu	Ala	Ala	Asp
		435					440					445			
Val	Ala	Lys	Ser	Gln	Phe	Leu	Ala	Thr	Val	Ser	His	Glu	Ile	Arg	Thr
450						455					460				
Pro	Met	Asn	Gly	Val	Leu	Gly	Met	Leu	His	Met	Leu	Met	Asp	Thr	Glu
465					470					475					480
Leu	Asp	Val	Thr	Gln	Gln	Asp	Tyr	Val	Arg	Thr	Ala	Gln	Ala	Ser	Gly
				485					490						495
Lys	Ala	Leu	Val	Ser	Leu	Ile	Asn	Glu	Val	Leu	Asp	Gln	Ala	Lys	Ile
			500					505					510		
Glu	Ser	Gly	Lys	Leu	Glu	Leu	Glu	Glu	Val	Arg	Phe	Asp	Leu	Arg	Gly
		515					520					525			
Ile	Leu	Asp	Asp	Val	Leu	Ser	Leu	Phe	Ser	Ser	Lys	Ser	Gln	Gln	Lys
530						535					540				
Gly	Val	Glu	Leu	Ala	Val	Tyr	Ile	Ser	Asp	Arg	Val	Pro	Asp	Met	Leu
545					550					555					560
Ile	Gly	Asp	Pro	Gly	Arg	Phe	Arg	Gln	Ile	Leu	Thr	Asn	Leu	Met	Gly
				565					570						575
Asn	Ser	Ile	Lys	Phe	Thr	Glu	Lys	Gly	His	Ile	Phe	Val	Thr	Val	His
			580					585					590		
Leu	Val	Asp	Glu	Leu	Phe	Glu	Ser	Ile	Asp	Gly	Glu	Thr	Ala	Ser	Ser
		595					600					605			
Pro	Glu	Ser	Thr	Leu	Ser	Gly	Leu	Pro	Val	Ala	Asp	Arg	Gln	Arg	Ser
610						615					620				
Trp	Glu	Asn	Phe	Lys	Ala	Phe	Ser	Ser	Asn	Gly	His	Arg	Ser	Phe	Glu
625					630					635					640
Pro	Ser	Pro	Pro	Asp	Ile	Asn	Leu	Ile	Val	Ser	Val	Glu	Asp	Thr	Gly
				645					650						655
Val	Gly	Ile	Pro	Val	Glu	Ala	Gln	Ser	Arg	Ile	Phe	Thr	Pro	Phe	Met
			660					665					670		
Gln	Val	Gly	Pro	Ser	Ile	Ser	Arg	Thr	His	Gly	Gly	Thr	Gly	Ile	Gly
		675					680						685		
Leu	Ser	Ile	Ser	Lys	Cys	Leu	Val	Gly	Leu	Met	Lys	Gly	Glu	Ile	Gly
690						695					700				
Phe	Ser	Ser	Thr	Pro	Lys	Val	Gly	Ser	Thr	Phe	Thr	Phe	Thr	Ala	Val
705					710					715					720
Phe	Ser	Asn	Gly	Met	Gln	Pro	Ala	Glu	Arg	Lys	Asn	Asp	Asn	Asn	Gln
				725					730						735
Pro	Ile	Phe	Ser	Glu	Phe	Arg	Gly	Met	Lys	Ala	Val	Val	Val	Asp	His
			740				745						750		
Arg	Pro	Ala	Arg	Ala	Lys	Val	Ser	Trp	Tyr	His	Phe	Gln	Arg	Leu	Gly
		755					760						765		
Ile	Arg	Val	Glu	Val	Val	Pro	Arg	Val	Glu	Gln	Ala	Leu	His	Tyr	Leu
770						775					780				
Lys	Ile	Gly	Thr	Thr	Thr	Val	Asn	Met	Ile	Leu	Ile	Glu	Gln	Glu	Ile
785					790					795					800
Trp	Asn	Arg	Glu	Ala	Asp	Asp	Phe	Ile	Lys	Lys	Leu	Gln	Lys	Asp	Pro
				805					810						815
Leu	Phe	Leu	Ser	Pro	Lys	Leu	Ile	Leu	Leu	Ala	Asn	Ser	Val	Glu	Ser
			820					825					830		

```

Ser Ile Ser Glu Ala Leu Cys Thr Gly Ile Asp Pro Pro Ile Val Ile
      835      840      845
Val Lys Pro Leu Arg Ala Ser Met Leu Ala Ala Thr Leu Gln Arg Gly
      850      855      860
Leu Gly Ile Gly Ile Arg Glu Pro Pro Gln His Lys Gly Pro Pro Ala
865      870      875      880
Leu Ile Leu Arg Asn Leu Leu Leu Gly Arg Lys Ile Leu Ile Val Asp
      885      890      895
Asp Asn Asn Val Asn Leu Arg Val Ala Ala Gly Ala Leu Lys Lys Tyr
      900      905      910
Gly Ala Asp Val Val Cys Ala Glu Ser Gly Ile Lys Ala Ile Ser Leu
      915      920      925
Leu Lys Pro Pro His Glu Phe Asp Ala Cys Phe Met Asp Ile Gln Met
      930      935      940
Pro Glu Met Asp Gly Phe Glu Ala Thr Arg Arg Ile Arg Asp Met Glu
945      950      955      960
Glu Glu Met Asn Lys Arg Ile Lys Asn Gly Glu Ala Leu Ile Val Glu
      965      970      975
Asn Gly Asn Lys Thr Ser Trp His Leu Pro Val Leu Ala Met Thr Ala
      980      985      990
Asp Val Ile Gln Ala Thr His Glu Glu Cys Leu Lys Cys Gly Met Asp
      995      1000      1005
Gly Tyr Val Ser Lys Pro Phe Glu Ala Glu Gln Leu Tyr Arg Glu Val
1010      1015      1020
Ser Arg Phe Phe Asn Ser Pro Ser Asp Thr Glu Ser
1025      1030      1035

```

<210> 28

<211> 1057

<212> PRT

<213> Arabidopsis thaliana

<400> 28

```

Met Asn Trp Ala Leu Asn Asn His Gln Glu Glu Glu Glu Glu Pro Arg
 1      5      10      15
Arg Ile Glu Ile Ser Asp Ser Glu Ser Leu Glu Asn Leu Lys Ser Ser
      20      25      30
Asp Phe Tyr Gln Leu Gly Gly Gly Gly Ala Leu Asn Ser Ser Glu Lys
      35      40      45
Pro Arg Lys Ile Asp Phe Trp Arg Ser Gly Leu Met Gly Phe Ala Lys
      50      55      60
Met Gln Gln Gln Gln Gln Leu Gln His Ser Val Ala Val Lys Met Asn
65      70      75      80
Asn Asn Asn Asn Asn Asp Leu Met Gly Asn Lys Lys Gly Ser Thr Phe
      85      90      95
Ile Gln Glu His Arg Ala Leu Leu Pro Lys Ala Leu Ile Leu Trp Ile
      100      105      110
Ile Ile Val Gly Phe Ile Ser Ser Gly Ile Tyr Gln Trp Met Asp Asp
      115      120      125
Ala Asn Lys Ile Arg Arg Glu Glu Val Leu Val Ser Met Cys Asp Gln
      130      135      140
Arg Ala Arg Met Leu Gln Asp Gln Phe Ser Val Ser Val Asn His Val
145      150      155      160
His Ala Leu Ala Ile Leu Val Ser Thr Phe His Tyr His Lys Asn Pro
      165      170      175
Ser Ala Ile Asp Gln Glu Thr Phe Ala Glu Tyr Thr Ala Arg Thr Ala
      180      185      190

```

Phe	Glu	Arg	Pro	Leu	Leu	Ser	Gly	Val	Ala	Tyr	Ala	Glu	Lys	Val	Val
		195					200					205			
Asn	Phe	Glu	Arg	Glu	Met	Phe	Glu	Arg	Gln	His	Asn	Trp	Val	Ile	Lys
	210					215					220				
Thr	Met	Asp	Arg	Gly	Glu	Pro	Ser	Pro	Val	Arg	Asp	Glu	Tyr	Ala	Pro
225					230					235					240
Val	Ile	Phe	Ser	Gln	Asp	Ser	Val	Ser	Tyr	Leu	Glu	Ser	Leu	Asp	Met
				245					250					255	
Met	Ser	Gly	Glu	Glu	Asp	Arg	Glu	Asn	Ile	Leu	Arg	Ala	Arg	Glu	Thr
			260					265					270		
Gly	Lys	Ala	Val	Leu	Thr	Ser	Pro	Phe	Arg	Leu	Leu	Glu	Thr	His	His
	275						280					285			
Leu	Gly	Val	Val	Leu	Thr	Phe	Pro	Val	Tyr	Lys	Ser	Ser	Leu	Pro	Glu
	290					295					300				
Asn	Pro	Thr	Val	Glu	Glu	Arg	Ile	Ala	Ala	Thr	Ala	Gly	Tyr	Leu	Gly
305					310					315					320
Gly	Ala	Phe	Asp	Val	Glu	Ser	Leu	Val	Glu	Asn	Leu	Leu	Gly	Gln	Leu
				325					330					335	
Ala	Gly	Asn	Gln	Ala	Ile	Val	Val	His	Val	Tyr	Asp	Ile	Thr	Asn	Ala
			340					345					350		
Ser	Asp	Pro	Leu	Val	Met	Tyr	Gly	Asn	Gln	Asp	Glu	Glu	Ala	Asp	Arg
	355						360					365			
Ser	Leu	Ser	His	Glu	Ser	Lys	Leu	Asp	Phe	Gly	Asp	Pro	Phe	Arg	Lys
	370					375					380				
His	Lys	Met	Ile	Cys	Arg	Tyr	His	Gln	Lys	Ala	Pro	Ile	Pro	Leu	Asn
385					390					395					400
Val	Leu	Thr	Thr	Val	Pro	Leu	Phe	Phe	Ala	Ile	Gly	Phe	Leu	Val	Gly
				405					410					415	
Tyr	Ile	Leu	Tyr	Gly	Ala	Ala	Met	His	Ile	Val	Lys	Val	Glu	Asp	Asp
			420					425					430		
Phe	His	Glu	Met	Gln	Glu	Leu	Lys	Val	Arg	Ala	Glu	Ala	Ala	Asp	Val
		435					440					445			
Ala	Lys	Ser	Gln	Phe	Leu	Ala	Thr	Val	Ser	His	Glu	Ile	Arg	Thr	Pro
	450					455					460				
Met	Asn	Gly	Ile	Leu	Gly	Met	Leu	Ala	Met	Leu	Leu	Asp	Thr	Glu	Leu
465					470				475						480
Ser	Ser	Thr	Gln	Arg	Asp	Tyr	Ala	Gln	Thr	Ala	Gln	Val	Cys	Gly	Lys
			485					490					495		
Ala	Leu	Ile	Ala	Leu	Ile	Asn	Glu	Val	Leu	Asp	Arg	Ala	Lys	Ile	Glu
			500					505					510		
Ala	Gly	Lys	Leu	Glu	Leu	Glu	Ser	Val	Pro	Phe	Asp	Ile	Arg	Ser	Ile
	515						520					525			
Leu	Asp	Asp	Val	Leu	Ser	Leu	Phe	Ser	Glu	Glu	Ser	Arg	Asn	Lys	Ser
	530					535					540				
Ile	Glu	Leu	Ala	Val	Phe	Val	Ser	Asp	Lys	Val	Pro	Glu	Ile	Val	Lys
545					550					555					560
Gly	Asp	Ser	Gly	Arg	Phe	Arg	Gln	Ile	Ile	Ile	Asn	Leu	Val	Gly	Asn
				565				570						575	
Ser	Val	Lys	Phe	Thr	Glu	Lys	Gly	His	Ile	Phe	Val	Lys	Val	His	Leu
			580					585					590		
Ala	Glu	Gln	Ser	Lys	Asp	Glu	Ser	Glu	Pro	Lys	Asn	Ala	Leu	Asn	Gly
	595					600					605				
Gly	Val	Ser	Glu	Glu	Met	Ile	Val	Val	Ser	Lys	Gln	Ser	Ser	Tyr	Asn
	610					615					620				
Thr	Leu	Ser	Gly	Tyr	Glu	Ala	Ala	Asp	Gly	Arg	Asn	Ser	Trp	Asp	Ser
625					630					635					640
Phe	Lys	His	Leu	Val	Ser	Glu	Glu	Gln	Ser	Leu	Ser	Glu	Phe	Asp	Ile
				645				650						655	

Ser Ser Asn Val Arg Leu Met Val Ser Ile Glu Asp Thr Gly Ile Gly
 660 665 670
 Ile Pro Leu Val Ala Gln Gly Arg Val Phe Met Pro Phe Met Gln Ala
 675 680 685
 Asp Ser Ser Thr Ser Arg Asn Tyr Gly Gly Thr Gly Ile Gly Leu Ser
 690 695 700
 Ile Ser Lys Cys Leu Val Glu Leu Met Arg Gly Gln Ile Asn Phe Ile
 705 710 715 720
 Ser Arg Pro His Ile Gly Ser Thr Phe Trp Phe Thr Ala Val Leu Glu
 725 730 735
 Lys Cys Asp Lys Cys Ser Ala Ile Asn His Met Lys Lys Pro Asn Val
 740 745 750
 Glu His Leu Pro Ser Thr Phe Lys Gly Met Lys Ala Ile Val Val Asp
 755 760 765
 Ala Lys Pro Val Arg Ala Ala Val Thr Arg Tyr His Met Lys Arg Leu
 770 775 780
 Gly Ile Asn Val Asp Val Val Thr Ser Leu Lys Thr Ala Val Val Ala
 785 790 795 800
 Ala Ala Ala Phe Glu Arg Asn Gly Ser Pro Leu Pro Thr Lys Pro Gln
 805 810 815
 Leu Asp Met Ile Leu Val Glu Lys Asp Ser Trp Ile Ser Thr Glu Asp
 820 825 830
 Asn Asp Ser Glu Ile Arg Leu Leu Asn Ser Arg Thr Asn Gly Asn Val
 835 840 845
 His His Lys Ser Pro Lys Leu Ala Leu Phe Ala Thr Asn Ile Thr Asn
 850 855 860
 Ser Glu Phe Asp Arg Ala Lys Ser Ala Gly Phe Ala Asp Thr Val Ile
 865 870 875 880
 Met Lys Pro Leu Arg Ala Ser Met Ile Gly Ala Cys Leu Gln Gln Val
 885 890 895
 Leu Glu Leu Arg Lys Thr Arg Gln Gln His Pro Glu Gly Ser Ser Pro
 900 905 910
 Ala Thr Leu Lys Ser Leu Leu Thr Gly Lys Lys Ile Leu Val Val Asp
 915 920 925
 Asp Asn Ile Val Asn Arg Arg Val Ala Ala Gly Ala Leu Lys Lys Phe
 930 935 940
 Gly Ala Glu Val Val Cys Ala Glu Ser Gly Gln Val Ala Leu Gly Leu
 945 950 955 960
 Leu Gln Ile Pro His Thr Phe Asp Ala Cys Phe Met Asp Ile Gln Met
 965 970 975
 Pro Gln Met Asp Gly Phe Glu Ala Thr Arg Gln Ile Arg Met Met Glu
 980 985 990
 Lys Glu Thr Lys Glu Lys Thr Asn Leu Glu Trp His Leu Pro Ile Leu
 995 1000 1005
 Ala Met Thr Ala Asp Val Ile His Ala Thr Tyr Glu Glu Cys Leu Lys
 1010 1015 1020
 Ser Gly Met Asp Gly Tyr Val Ser Lys Pro Phe Glu Glu Glu Asn Leu
 1025 1030 1035 1040
 Tyr Lys Ser Val Ala Lys Ser Phe Lys Pro Asn Pro Ile Ser Pro Ser
 1045 1050 1055
 Ser

<210> 29

<211> 1057

<212> PRT

<213> Artificial Sequence

<220>

<223> derived from *Arabidopsis thaliana*

<400> 29

Met	Asn	Trp	Ala	Leu	Asn	Asn	His	Gln	Glu	Glu	Glu	Glu	Glu	Pro	Arg
1				5					10					15	
Arg	Ile	Glu	Ile	Ser	Asp	Ser	Glu	Ser	Leu	Glu	Asn	Leu	Lys	Ser	Ser
			20					25					30		
Asp	Phe	Tyr	Gln	Leu	Gly	Gly	Gly	Gly	Ala	Leu	Asn	Ser	Ser	Glu	Lys
		35					40					45			
Pro	Arg	Lys	Ile	Asp	Phe	Trp	Arg	Ser	Gly	Leu	Met	Gly	Phe	Ala	Lys
	50					55					60				
Met	Gln	Gln	Gln	Gln	Gln	Gln	Leu	Gln	His	Ser	Val	Ala	Val	Lys	Met
65					70						75				80
Asn	Asn	Asn	Asn	Asn	Asp	Leu	Met	Gly	Asn	Lys	Lys	Gly	Ser	Thr	Phe
				85					90					95	
Ile	Gln	Glu	His	Arg	Ala	Leu	Leu	Pro	Lys	Ala	Leu	Ile	Leu	Trp	Ile
			100					105					110		
Ile	Ile	Val	Gly	Phe	Ile	Ser	Ser	Gly	Ile	Tyr	Gln	Trp	Met	Asp	Asp
		115					120					125			
Ala	Asn	Lys	Ile	Arg	Arg	Glu	Glu	Val	Leu	Val	Ser	Met	Cys	Asp	Gln
	130					135					140				
Arg	Ala	Arg	Met	Leu	Gln	Asp	Gln	Phe	Ser	Val	Ser	Val	Asn	His	Val
145					150					155					160
His	Ala	Leu	Ala	Ile	Leu	Val	Ser	Thr	Phe	His	Tyr	His	Lys	Asn	Pro
			165						170					175	
Ser	Ala	Ile	Asp	Gln	Glu	Thr	Phe	Ala	Glu	Tyr	Thr	Ala	Arg	Thr	Ala
		180					185						190		
Phe	Glu	Arg	Pro	Leu	Leu	Ser	Gly	Val	Ala	Tyr	Ala	Glu	Lys	Val	Val
	195					200						205			
Asn	Phe	Glu	Arg	Glu	Met	Phe	Glu	Arg	Gln	His	Asn	Trp	Val	Ile	Lys
	210					215					220				
Thr	Met	Asp	Arg	Gly	Glu	Pro	Ser	Pro	Val	Arg	Asp	Glu	Tyr	Ala	Pro
225					230					235					240
Val	Ile	Phe	Ser	Gln	Asp	Ser	Val	Ser	Tyr	Leu	Glu	Ser	Leu	Asp	Met
			245						250					255	
Met	Ser	Gly	Glu	Glu	Asp	Arg	Glu	Asn	Ile	Leu	Arg	Ala	Arg	Glu	Thr
		260					265						270		
Gly	Lys	Ala	Val	Leu	Thr	Ser	Pro	Phe	Arg	Leu	Leu	Glu	Thr	His	His
		275					280					285			
Leu	Gly	Val	Val	Leu	Thr	Phe	Pro	Val	Tyr	Lys	Ser	Ser	Leu	Pro	Glu
	290					295					300				
Asn	Pro	Thr	Val	Glu	Glu	Arg	Ile	Ala	Ala	Thr	Ala	Gly	Tyr	Leu	Gly
305					310					315					320
Gly	Ala	Phe	Asp	Val	Glu	Ser	Leu	Val	Glu	Asn	Leu	Leu	Gly	Gln	Leu
			325						330					335	
Ala	Gly	Asn	Gln	Ala	Ile	Val	Val	His	Val	Tyr	Asp	Ile	Thr	Asn	Ala
		340						345					350		
Ser	Asp	Pro	Leu	Val	Met	Tyr	Gly	Asn	Gln	Asp	Glu	Glu	Ala	Asp	Arg
	355					360					365				
Ser	Leu	Ser	His	Glu	Ser	Lys	Leu	Asp	Phe	Gly	Asp	Pro	Phe	Arg	Lys
	370					375					380				
His	Lys	Met	Ile	Cys	Arg	Tyr	His	Gln	Lys	Ala	Pro	Ile	Pro	Leu	Asn
385					390					395					400
Val	Leu	Thr	Thr	Val	Pro	Leu	Phe	Phe	Ala	Ile	Gly	Phe	Leu	Val	Gly
			405						410					415	
Tyr	Ile	Leu	Tyr	Gly	Ala	Ala	Met	His	Ile	Val	Lys	Val	Glu	Asp	Asp
			420					425					430		

Phe	His	Glu	Met	Gln	Glu	Leu	Lys	Val	Arg	Ala	Glu	Ala	Ala	Asp	Val
		435					440					445			
Ala	Lys	Ser	Gln	Phe	Leu	Ala	Thr	Val	Ser	Gln	Glu	Ile	Arg	Thr	Pro
	450					455					460				
Met	Asn	Gly	Ile	Leu	Gly	Met	Leu	Ala	Met	Leu	Leu	Asp	Thr	Glu	Leu
465					470					475					480
Ser	Ser	Thr	Gln	Arg	Asp	Tyr	Ala	Gln	Thr	Ala	Gln	Val	Cys	Gly	Lys
			485					490					495		
Ala	Leu	Ile	Ala	Leu	Ile	Asn	Glu	Val	Leu	Asp	Arg	Ala	Lys	Ile	Glu
			500					505					510		
Ala	Gly	Lys	Leu	Glu	Leu	Glu	Ser	Val	Pro	Phe	Asp	Ile	Arg	Ser	Ile
	515						520					525			
Leu	Asp	Asp	Val	Leu	Ser	Leu	Phe	Ser	Glu	Glu	Ser	Arg	Asn	Lys	Ser
	530					535					540				
Ile	Glu	Leu	Ala	Val	Phe	Val	Ser	Asp	Lys	Val	Pro	Glu	Ile	Val	Lys
545					550					555					560
Gly	Asp	Ser	Gly	Arg	Phe	Arg	Gln	Ile	Ile	Ile	Asn	Leu	Val	Gly	Asn
			565					570						575	
Ser	Val	Lys	Phe	Thr	Glu	Lys	Gly	His	Ile	Phe	Val	Lys	Val	His	Leu
			580					585					590		
Ala	Glu	Gln	Ser	Lys	Asp	Glu	Ser	Glu	Pro	Lys	Asn	Ala	Leu	Asn	Gly
		595					600					605			
Gly	Val	Ser	Glu	Glu	Met	Ile	Val	Val	Ser	Lys	Gln	Ser	Ser	Tyr	Asn
	610					615					620				
Thr	Leu	Ser	Gly	Tyr	Glu	Ala	Ala	Asp	Gly	Arg	Asn	Ser	Trp	Asp	Ser
625					630					635					640
Phe	Lys	His	Leu	Val	Ser	Glu	Glu	Gln	Ser	Leu	Ser	Glu	Phe	Asp	Ile
			645					650						655	
Ser	Ser	Asn	Val	Arg	Leu	Met	Val	Ser	Ile	Glu	Asp	Thr	Gly	Ile	Gly
		660					665						670		
Ile	Pro	Leu	Val	Ala	Gln	Gly	Arg	Val	Phe	Met	Pro	Phe	Met	Gln	Ala
	675					680						685			
Asp	Ser	Ser	Thr	Ser	Arg	Asn	Tyr	Gly	Gly	Thr	Gly	Ile	Gly	Leu	Ser
690					695						700				
Ile	Ser	Lys	Cys	Leu	Val	Glu	Leu	Met	Arg	Gly	Gln	Ile	Asn	Phe	Ile
705					710					715					720
Ser	Arg	Pro	His	Ile	Gly	Ser	Thr	Phe	Trp	Phe	Thr	Ala	Val	Leu	Glu
			725					730						735	
Lys	Cys	Asp	Lys	Cys	Ser	Ala	Ile	Asn	His	Met	Lys	Lys	Pro	Asn	Val
		740						745					750		
Glu	His	Leu	Pro	Ser	Thr	Phe	Lys	Gly	Met	Lys	Ala	Ile	Val	Val	Asp
	755						760					765			
Ala	Lys	Pro	Val	Arg	Ala	Ala	Val	Thr	Arg	Tyr	His	Met	Lys	Arg	Leu
	770					775					780				
Gly	Ile	Asn	Val	Asp	Val	Val	Thr	Ser	Leu	Lys	Thr	Ala	Val	Val	Ala
785					790					795					800
Ala	Ala	Ala	Phe	Glu	Arg	Asn	Gly	Ser	Pro	Leu	Pro	Thr	Lys	Pro	Gln
			805					810						815	
Leu	Asp	Met	Ile	Leu	Val	Glu	Lys	Asp	Ser	Trp	Ile	Ser	Thr	Glu	Asp
		820						825					830		
Asn	Asp	Ser	Glu	Ile	Arg	Leu	Leu	Asn	Ser	Arg	Thr	Asn	Gly	Asn	Val
	835						840						845		
His	His	Lys	Ser	Pro	Lys	Leu	Ala	Leu	Phe	Ala	Thr	Asn	Ile	Thr	Asn
	850					855					860				
Ser	Glu	Phe	Asp	Arg	Ala	Lys	Ser	Ala	Gly	Phe	Ala	Asp	Thr	Val	Ile
865					870					875					880
Met	Lys	Pro	Leu	Arg	Ala	Ser	Met	Ile	Gly	Ala	Cys	Leu	Gln	Gln	Val
			885						890					895	

```

Leu Glu Leu Arg Lys Thr Arg Gln Gln His Pro Glu Gly Ser S r Pro
      900      905      910
Ala Thr Leu Lys Ser Leu Leu Thr Gly Lys Lys Ile Leu Val Val Asp
      915      920      925
Asp Asn Ile Val Asn Arg Arg Val Ala Ala Gly Ala Leu Lys Lys Phe
      930      935      940
Gly Ala Glu Val Val Cys Ala Glu Ser Gly Gln Val Ala Leu Gly Leu
945      950      955      960
Leu Gln Ile Pro His Thr Phe Asp Ala Cys Phe Met Asp Ile Gln Met
      965      970      975
Pro Gln Met Asp Gly Phe Glu Ala Thr Arg Gln Ile Arg Met Met Glu
      980      985      990
Lys Glu Thr Lys Glu Lys Thr Asn Leu Glu Trp His Leu Pro Ile Leu
      995      1000      1005
Ala Met Thr Ala Asp Val Ile His Ala Thr Tyr Glu Glu Cys Leu Lys
      1010      1015      1020
Ser Gly Met Asp Gly Tyr Val Ser Lys Pro Phe Glu Glu Glu Asn Leu
1025      1030      1035      1040
Tyr Lys Ser Val Ala Lys Ser Phe Lys Pro Asn Pro Ile Ser Pro Ser
      1045      1050      1055
Ser

```

```

<210> 30
<211> 1057
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> derived from Arabidopsis thaliana

```

```

<400> 30
Met Asn Trp Ala Leu Asn Asn His Gln Glu Glu Glu Glu Glu Pro Arg
 1      5      10      15
Arg Ile Glu Ile Ser Asp Ser Glu Ser Leu Glu Asn Leu Lys Ser Ser
      20      25      30
Asp Phe Tyr Gln Leu Gly Gly Gly Gly Ala Leu Asn Ser Ser Glu Lys
      35      40      45
Pro Arg Lys Ile Asp Phe Trp Arg Ser Gly Leu Met Gly Phe Ala Lys
      50      55      60
Met Gln Gln Gln Gln Gln Leu Gln His Ser Val Ala Val Lys Met Asn
65      70      75      80
Asn Asn Asn Asn Asn Asp Leu Met Gly Asn Lys Lys Gly Ser Thr Phe
      85      90      95
Ile Gln Glu His Arg Ala Leu Leu Pro Lys Ala Leu Ile Leu Trp Ile
      100      105      110
Ile Ile Val Gly Phe Ile Ser Ser Gly Ile Tyr Gln Trp Met Asp Asp
      115      120      125
Ala Asn Lys Ile Arg Arg Glu Glu Val Leu Val Ser Met Cys Asp Gln
      130      135      140
Arg Ala Arg Met Leu Gln Asp Gln Phe Ser Val Ser Val Asn His Val
145      150      155      160
His Ala Leu Ala Ile Leu Val Ser Thr Phe His Tyr His Lys Asn Pro
      165      170      175
Ser Ala Ile Asp Gln Glu Thr Phe Ala Glu Tyr Thr Ala Arg Thr Ala
      180      185      190
Phe Glu Arg Pro Leu Leu Ser Gly Val Ala Tyr Ala Glu Lys Val Val

```

		195				200				205					
Asn	Phe	Glu	Arg	Glu	Met	Phe	Glu	Arg	Gln	His	Asn	Trp	Val	Ile	Lys
210						215					220				
Thr	Met	Asp	Arg	Gly	Glu	Pro	Ser	Pro	Val	Arg	Asp	Glu	Tyr	Ala	Pro
225					230					235					240
Val	Ile	Phe	Ser	Gln	Asp	Ser	Val	Ser	Tyr	Leu	Glu	Ser	Leu	Asp	Met
				245					250					255	
Met	Ser	Gly	Glu	Glu	Asp	Arg	Glu	Asn	Ile	Leu	Arg	Ala	Arg	Glu	Thr
			260					265					270		
Gly	Lys	Ala	Val	Leu	Thr	Ser	Pro	Phe	Arg	Leu	Leu	Glu	Thr	His	His
		275					280					285			
Leu	Gly	Val	Val	Leu	Thr	Phe	Pro	Val	Tyr	Lys	Ser	Ser	Leu	Pro	Glu
290						295					300				
Asn	Pro	Thr	Val	Glu	Glu	Arg	Ile	Ala	Ala	Thr	Ala	Gly	Tyr	Leu	Gly
305					310					315					320
Gly	Ala	Phe	Asp	Val	Glu	Ser	Leu	Val	Glu	Asn	Leu	Leu	Gly	Gln	Leu
				325					330					335	
Ala	Gly	Asn	Gln	Ala	Ile	Val	Val	His	Val	Tyr	Asp	Ile	Thr	Asn	Ala
			340					345					350		
Ser	Asp	Pro	Leu	Val	Met	Tyr	Gly	Asn	Gln	Asp	Glu	Glu	Ala	Asp	Arg
		355					360					365			
Ser	Leu	Ser	His	Glu	Ser	Lys	Leu	Asp	Phe	Gly	Asp	Pro	Phe	Arg	Lys
370						375					380				
His	Lys	Met	Ile	Cys	Arg	Tyr	His	Gln	Lys	Ala	Pro	Ile	Pro	Leu	Asn
385					390					395					400
Val	Leu	Thr	Thr	Val	Pro	Leu	Phe	Phe	Ala	Ile	Gly	Phe	Leu	Val	Gly
				405					410					415	
Tyr	Ile	Leu	Tyr	Gly	Ala	Ala	Met	His	Ile	Val	Lys	Val	Glu	Asp	Asp
			420					425					430		
Phe	His	Glu	Met	Gln	Glu	Leu	Lys	Val	Arg	Ala	Glu	Ala	Ala	Asp	Val
		435					440					445			
Ala	Lys	Ser	Gln	Phe	Leu	Ala	Thr	Val	Ser	His	Glu	Ile	Arg	Thr	Pro
450						455					460				
Met	Asn	Gly	Ile	Leu	Gly	Met	Leu	Ala	Met	Leu	Leu	Asp	Thr	Glu	Leu
465					470					475					480
Ser	Ser	Thr	Gln	Arg	Asp	Tyr	Ala	Gln	Thr	Ala	Gln	Val	Cys	Gly	Lys
				485					490					495	
Ala	Leu	Ile	Ala	Leu	Ile	Asn	Glu	Val	Leu	Asp	Arg	Ala	Lys	Ile	Glu
			500					505					510		
Ala	Gly	Lys	Leu	Glu	Leu	Glu	Ser	Val	Pro	Phe	Asp	Ile	Arg	Ser	Ile
		515					520					525			
Leu	Asp	Asp	Val	Leu	Ser	Leu	Phe	Ser	Glu	Glu	Ser	Arg	Asn	Lys	Ser
530						535					540				
Ile	Glu	Leu	Ala	Val	Phe	Val	Ser	Asp	Lys	Val	Pro	Glu	Ile	Val	Lys
545					550					555					560
Gly	Asp	Ser	Gly	Arg	Phe	Arg	Gln	Ile	Ile	Ile	Asn	Leu	Val	Gly	Asn
				565					570					575	
Ser	Val	Lys	Phe	Thr	Glu	Lys	Gly	His	Ile	Phe	Val	Lys	Val	His	Leu
			580					585					590		
Ala	Glu	Gln	Ser	Lys	Asp	Glu	Ser	Glu	Pro	Lys	Asn	Ala	Leu	Asn	Gly
		595					600					605			
Gly	Val	Ser	Glu	Glu	Met	Ile	Val	Val	Ser	Lys	Gln	Ser	Ser	Tyr	Asn
610						615					620				
Thr	Leu	Ser	Gly	Tyr	Glu	Ala	Ala	Asp	Gly	Arg	Asn	Ser	Trp	Asp	Ser
625					630					635					640
Phe	Lys	His	Leu	Val	Ser	Glu	Glu	Gln	Ser	Leu	Ser	Glu	Phe	Asp	Ile
				645					650					655	
Ser	Ser	Asn	Val	Arg	Leu	Met	Val	Ser	Ile	Glu	Asp	Thr	Gly	Ile	Gly

			660					665					670			
Ile	Pro	Leu	Val	Ala	Gln	Gly	Arg	Val	Phe	Met	Pro	Phe	Met	Gln	Ala	
		675					680					685				
Asp	Ser	Ser	Thr	Ser	Arg	Asn	Tyr	Gly	Gly	Thr	Gly	Ile	Gly	Leu	Ser	
	690					695					700					
Ile	Ser	Lys	Cys	Leu	Val	Glu	Leu	Met	Arg	Gly	Gln	Ile	Asn	Phe	Ile	
705					710					715					720	
Ser	Arg	Pro	His	Ile	Gly	Ser	Thr	Phe	Trp	Phe	Thr	Ala	Val	Leu	Glu	
				725					730					735		
Lys	Cys	Asp	Lys	Cys	Ser	Ala	Ile	Asn	His	Met	Lys	Lys	Pro	Asn	Val	
		740						745					750			
Glu	His	Leu	Pro	Ser	Thr	Phe	Lys	Gly	Met	Lys	Ala	Ile	Val	Val	Asp	
	755						760					765				
Ala	Lys	Pro	Val	Arg	Ala	Ala	Val	Thr	Arg	Tyr	His	Met	Lys	Arg	Leu	
	770					775					780					
Gly	Ile	Asn	Val	Asp	Val	Val	Thr	Ser	Leu	Lys	Thr	Ala	Val	Val	Ala	
785					790					795					800	
Ala	Ala	Ala	Phe	Glu	Arg	Asn	Gly	Ser	Pro	Leu	Pro	Thr	Lys	Pro	Gln	
				805					810					815		
Leu	Asp	Met	Ile	Leu	Val	Glu	Lys	Asp	Ser	Trp	Ile	Ser	Thr	Glu	Asp	
		820						825					830			
Asn	Asp	Ser	Glu	Ile	Arg	Leu	Leu	Asn	Ser	Arg	Thr	Asn	Gly	Asn	Val	
	835						840					845				
His	His	Lys	Ser	Pro	Lys	Leu	Ala	Leu	Phe	Ala	Thr	Asn	Ile	Thr	Asn	
	850					855					860					
Ser	Glu	Phe	Asp	Arg	Ala	Lys	Ser	Ala	Gly	Phe	Ala	Asp	Thr	Val	Ile	
865					870					875					880	
Met	Lys	Pro	Leu	Arg	Ala	Ser	Met	Ile	Gly	Ala	Cys	Leu	Gln	Gln	Val	
				885					890					895		
Leu	Glu	Leu	Arg	Lys	Thr	Arg	Gln	Gln	His	Pro	Glu	Gly	Ser	Ser	Pro	
		900						905					910			
Ala	Thr	Leu	Lys	Ser	Leu	Leu	Thr	Gly	Lys	Lys	Ile	Leu	Val	Val	Asp	
	915						920					925				
Asp	Asn	Ile	Val	Asn	Arg	Arg	Val	Ala	Ala	Gly	Ala	Leu	Lys	Lys	Phe	
	930					935					940					
Gly	Ala	Glu	Val	Val	Cys	Ala	Glu	Ser	Gly	Gln	Val	Ala	Leu	Gly	Leu	
945					950					955					960	
Leu	Gln	Ile	Pro	His	Thr	Phe	Asp	Ala	Cys	Phe	Met	Asn	Ile	Gln	Met	
				965					970					975		
Pro	Gln	Met	Asp	Gly	Phe	Glu	Ala	Thr	Arg	Gln	Ile	Arg	Met	Met	Glu	
		980						985					990			
Lys	Glu	Thr	Lys	Glu	Lys	Thr	Asn	Leu	Glu	Trp	His	Leu	Pro	Ile	Leu	
	995						10									